

Albizia lebbek Stem Bark Aqueous Extract as Alternative to Antibiotic Feed Additives in Broiler Chicks Diets: Haematology, Serum Indices and Oxidative Status

Alagbe, J.O.^{1*}, Ajagbe A.D.², Attama Jeremiah³, Philemon K.C.⁴ and Bello, Kamoru. A⁵

¹Department of Animal Nutrition and Biochemistry, Sumitra Research Institute, Gujarat, India.

²Department of Animal Production, Kogi State University, Anyigba, Kogi State, Nigeria.

³Department of Veterinary Medicine, University of Nigeria, Nsukka.

⁴Department of Theriogenology, University of Abuja, Nigeria.

⁵Al-Hikmah University Ilorin, Nigeria

***Corresponding Author:** Alagbe, J.O, 1Department of Animal Nutrition and Biochemistry, Sumitra Research Institute, Gujarat, India.

Received date: August 15, 2020; **Accepted date:** October 09, 2020; **Published date:** October 14, 2020

Citation: Alagbe, J.O, Ajagbe A.D, Attama J, Philemon K.C, Bello K. A. (2020) Albizia lebbek Stem Bark Aqueous Extract as Alternative to Antibiotic Feed Additives in Broiler Chicks Diets: Haematology, Serum Indices and Oxidative Status. *Biomedical Research and Clinical Reviews*. 1(3); DOI:10.31579/2692-9406/018

Copyright:©2020 Alagbe, J.O, This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Abstract

A total of Three hundred and seventy five (375) one day old (Ross 308) broiler chicks with mixed sex were used to examine the effects of Albizia lebbek stem bark (ATSM) aqueous extract as alternative to antibiotic feed additives in broiler chicks diets: haematology, serum biochemical indices and oxidative status. Birds were divided to five treatments with five replicates of fifteen (15) birds in a completely randomized design. Treatment 1 (basal diet + 0 % ATSM), treatment 2 (basal diet +1.2 grams Oxytetracycline per litre of water), treatment 3 (basal diet + 10 ml ATSM per liter of water), treatment 4 (basal diet + 20 ml ATSM per litre of water) and treatment 5 (basal diet + 30 ml ATSM per liter of water) and the trial lasted for 56 days. Results on some haematological parameters revealed that red blood cell (RBC), pack cell volume (PCV), haemoglobin (Hb), mean corpuscular volume (MCV), mean corpuscular haemoglobin (MCH), mean corpuscular haemoglobin concentration (MCHC), white blood cell (WBC) and its differentials were significantly ($P<0.05$) different among the treatments. Total protein, glucose, urea, cholesterol, creatinine, aspartate aminotransaminase (AST) and alanine aminotransferase (ALT) were significantly ($P<0.05$) affected by ATSM. Activities of superoxide dismutase (SDA), glutathione peroxidase (GPx), catalase (CAT) and malonyldialdehyde (MLA) were significantly influenced by ATSM ($P<0.05$). It was concluded that ATSM could be administered to broiler chicks at 30 ml/litre without any negative effect on the general performance of birds.

Key words: Albizia lebbek; broiler chicks; haematology; serum biochemical indices.

Introduction

Phytogenics are heterogeneous group of feed additives emanating from plants and consists of herbs, fruit, spices and other plant parts (Santi and Kim, 2017). According to Veerschari et al. (2011), there are over 100, 000 species of plants used globally for medicinal purposes, many have been used in the form of therapy for livestock among resource poor smallholder farmers to treat variety of conditions of animals (Mirazaei-Aghsaghali, 2012; Alagbe et al., 2020). Most medicinal plants have been found to be abundant in minerals, vitamins, amino acid and bioactive chemicals [phytochemicals] (Olafadehan et al., 2020). However, only a small percentage have been explored or studied for their pharmacological properties (WHO, 1992). Nutrients in plants have great influence on responses of animals to a disease challenge and it has a direct correlation to the immune system (Gary and Richard, 2002). One of the numerous plant used for therapeutic purposes is Albizia lebbek.

Albizia lebbek (Mimosaceae) is a perennial tree native to tropical and subtropical regions of Asia and Africa. The genus Albizia comprises of almost 150 species spread all over India, China, Nigeria, Senegal, Ghana,

Togo, Congo, Benin, Angola, Uganda, Botswana among others (Ukpabi and Offor, 2018; Karuppannan, 2013). The plant parts (stem, leaf and seeds) have been found to be loaded with minerals (calcium, phosphorus, iron, copper, zinc, selenium, molybdenum and potassium), vitamins and amino acids (Alagbe and Soares, 2018; Uzoekwe and Mohammed, 2015; Mohammed et al., 2012). The leaf and stem has traditionally been used for the treatment of fever, tooth ache, wounds, leprosy, ulcer, cold, leprosy, sexually transmitted diseases and other respiratory infections (Labaran, 2016; Uwaya et al., 2017).

Several reports on the biological activity of Albizia lebbek revealed that the plant performs antimicrobial (Labaran et al., 2016), anti-inflammatory (Gupta et al., 2004), antioxidant (Mc Donald et al., 2001), analgesic (Sharma et al., 2007), antihelminthic (Karuppannan, 2013), hepatoprotective (Edeoga et al., 2005; Alagbe, 2019), antidiabetic (Kareru et al., 2007), immuno-modulatory (Sharma et al., 2007) and antihyperlipidemic properties (Ueda et al., 2003) due to the presence of several bioactive chemicals such as alkaloids, flavonoids, saponins, phenols, tannin etc (Labaran et al., 2016). In view of these abundant potential, administration

of Albizia lebeck stem bark to birds will possibly supply nutrients to meet all the body's need during a time of challenge.

Therefore, this experiment was designed to determine the effects of Albizia lebeck stem bark aqueous extract as alternative to antibiotic feed additives in broiler chicks diets: haematology, serum biochemical indices and oxidative status.

Materials and Methods

Study Area

The experiment was carried out at Division of Animal Nutrition, Sumitra Research Institute, Gujarat, India during the month of April to June, 2019.

Sources, collection and preparation of Albizia lebeck stem bark extract

The stem of Albizia lebeck stems were obtained from different plants in Gujarat, India and authenticated by a botanist Sharma Xing. The stem bark were cut into pieces and thoroughly washed with distilled water, air dried under the shade to maintain the bioactive chemicals in the test material. The dried samples were pulverized into powder using pestle and mortar, thereafter 250 grams of the sample was soaked into 1000 litres of water, sample was continuously stirred and kept in the refrigerator at 4°C for 72 hours. All mixtures were filtered using Whatman filter paper and the filterates (ATSM) were collected into a clean labelled container.

Experimental animals and management

Three hundred and seventy five one day old (Ross 308) broiler chicks with mixed sex were used for the experiment. The birds were purchased

from a commercial hatchery in India and weighed on arrival on the farm to obtain their initial body weight and thereafter weekly. A deep litter housing system was used, it was fumigated two weeks prior to the commencement of the study, and the surrounding environment was also cleaned daily to ensure proper hygiene. Birds were divided to five treatments with five replicates of fifteen (15) birds in a completely randomized design. Electric brooders were used and wood shavings serve as the litter material. Daily feed intake (g) was calculated as a difference between feed offered and left-over. Vaccines were administered according to the prevailing disease condition in the environment and all other management practices were strictly adhered to throughout the experiment which lasted for 56 days.

Ration formulation

Three (3) basal diets were formulated at different stages of production to meet up with the requirements of birds according to NRC (1994). Broiler starter's mash (0-21 days), growers mash (22-35 days) and finishers mash (36-56 days).

Treatment 1 (basal diet + 0 % ATSM), treatment 2 (basal diet +1.2 grams Oxytetracycline per litre of water), treatment 3 (basal diet + 10 ml ATSM per liter of water), treatment 4 (basal diet + 20 ml ATSM per litre of water) and treatment 5 (basal diet + 30 ml ATSM per liter of water).

Parameters measured

Proximate compositions of experiment diet were determined by using official method of analysis by AOAC (2000). Amino acid analysis was carried out using amino acid analyzer with ion exchange chromatographic method (Model NH-09b) India.

Materials	Starter (1-21 days)	Grower (22-35 days)	Finisher (36-56 days)
Maize	50.00	56.00	60.50
Wheat offal	8.00	7.00	8.05
Soya meal	28.55	22.00	21.00
Groundnut cake	10.00	11.55	6.05
Fish meal	2.00	2.00	2.00
Bone meal	0.35	0.40	0.40
Limestone	0.20	0.20	0.20
Lysine	0.15	0.15	0.15
Methionine	0.20	0.20	0.20
Premix	0.25	0.25	0.25
Salt	0.30	0.30	0.30
TOTAL	100.0	100.0	100.0
Calculated analysis			
Crude protein	23.08	20.11	19.33
Ether extract	5.03	4.87	4.28
Crude fibre	3.06	3.95	3.42
Calcium	0.98	1.00	1.10
Phosphorus	0.47	0.40	0.51
Lysine	1.17	1.29	1.60
Meth +Cyst	0.87	0.82	0.51
ME (Kcal/kg)	2936	3000.8	3100.2

*Premix supplied per kg diet: - vit A, 13,000 IU; vit E, 5mg; vit D3, 3000IU, vit K, 3mg; vit B2, 5.5mg; Niacin, 25mg; vit B12, 16mg; choline chloride, 120mg; Mn, 5.2mg; Zn, 25mg; Cu, 2.6g; folic acid, 2mg; Fe, 5g; pantothenic acid, 10mg; biotin, 30.5g; antioxidant, 56mg.

Table 1: Chemical composition of experimental diets

Haematological and serum biochemical analysis

Blood samples were collected very early in the morning from the wing vein from three (3) randomly selected birds per replicate into a 5 ml sterile syringe using 23 gauge needles and transferred into an ethylene diamine tetra acetic acid (EDTA) bottle. Haematological parameters: pack cell

volume (PCV), red blood cell (RBC), haemoglobin (Hb), mean corpuscular haemoglobin (MCH), mean corpuscular haemoglobin concentration (MCHC), mean corpuscular volume (MCV), white blood cell (WBC) and its differentials were analyzed using an automated machine (Sysmex, Model KU-30 HG, India).

Serum analysis was carried out using bottles free from EDTA, blood were analyzed for total protein, albumin, globulin, glucose, cholesterol,

creatinine, alanine transaminase (ALT) and aspartate transaminase (AST) were assayed using diagnostic kit manufactured by Merck India Ltd (Model PS-09R) as described by Olubukola et al. (2015).

Antioxidant status

Activity of superoxide dismutase (SDA), glutathione peroxidase (GPx), catalase (CAT) and malonyldialdehyde (MLA) were carried out using

method outlined by Mahipal et al. (2015).

Statistical analysis

All data were subjected to one-way analysis of variance (ANOVA) using SPSS (18.0) and significant means were separated using Duncan multiple range tests (Duncan, 1955). Significant was declared if $P \leq 0.05$.

Amino acids	Composition (%)	*Reference level
Lysine	0.86	5.50
Arginine	1.65	1.00
Aspartic acid	2.00	-
Threonine	1.12	0.65
Histidine	3.45	0.30
Serine	0.78	-
Glycine	1.00	1.20
Alanine	3.00	-
Cystine	4.11	0.35
Valine	0.65	0.82
Leucine	1.00	1.20
Phenylalanine	0.34	0.50
Tyrosine	0.10	0.45
Isoleucine	2.00	0.60
Proline	0.03	0.20
Methionine	0.02	0.35

*NRC (1994)

Table 2: Amino acid compositions of Albizia lebbeck stem bark

Parameters	T1	T2	T3	T4	T5	SEM
PCV (%)	26.50 ^c	29.31 ^b	31.02 ^b	33.56 ^a	34.00 ^a	0.37
Hb (g/dl)	9.12 ^c	10.21 ^b	11.93 ^b	12.11 ^a	12.50 ^a	0.64
RBC $\times 10^6/\mu\text{l}$	1.88 ^c	2.00 ^b	2.10 ^b	2.60 ^a	2.97 ^a	0.07
MCV (fl)	111.2 ^b	119.8 ^a	120.5 ^a	123.6 ^a	130.4 ^a	8.10
MCH (pg)	34.51 ^c	50.43 ^b	56.11 ^b	57.67 ^a	59.00 ^a	2.51
MCHC (g/dl)	29.80 ^b	35.60 ^a	39.00 ^b	39.10 ^a	40.03 ^a	0.88
WBC $\times 10^3/\mu\text{l}$	20.41 ^b	20.62 ^b	22.74 ^b	22.88 ^b	30.04 ^a	0.12
Differentials ($10^3/\mu\text{l}$)						
Lymphocytes	10.45 ^c	14.08 ^b	15.44 ^b	18.71 ^b	20.04 ^a	1.96
Monocytes	0.07 ^c	1.11 ^b	1.20 ^b	1.26 ^b	1.72 ^a	0.01
Heterophils	4.23 ^b	5.06 ^a	5.40 ^a	5.89 ^a	6.11 ^a	0.41
Eosinophils	0.88 ^b	1.02 ^a	1.09 ^a	1.21 ^a	1.27 ^a	0.02

Means in the same row with different superscript are significantly different ($P < 0.05$)

Table 3: Haematological parameters of broiler chicks fed different levels ATSM

Parameters	T1	T2	T3	T4	T5	SEM
Total protein (g/dl)	2.57 ^b	3.22 ^a	3.69 ^a	3.88 ^a	3.97 ^a	0.67
Albumin (g/dl)	1.34 ^b	1.55 ^b	1.91 ^b	2.00 ^a	2.03 ^a	0.02
Globulin (g/dl)	1.23 ^c	1.67 ^b	1.78 ^b	1.88 ^a	1.94 ^a	0.15
Creatinine (mg/dl)	0.15 ^c	0.45 ^b	0.81 ^a	0.87 ^a	0.91 ^a	0.01
Glucose (mg/dl)	196.1 ^c	204.3 ^a	234.1 ^a	241.5 ^a	250.6 ^a	4.33
Cholesterol (mg/dl)	101.4 ^a	98.6 ^b	90.4 ^a	89.4 ^a	87.5 ^a	2.87
Uric acid (mg/dl)	7.33 ^a	4.89 ^b	4.22 ^b	4.00 ^b	3.88 ^b	0.05
ALT (u/l)	74.1 ^a	70.5 ^a	61.6 ^b	58.1 ^b	50.7 ^b	1.45
AST (u/l)	300.7 ^a	288.5 ^b	230.4 ^b	218.0 ^b	200.9 ^b	9.45

Means in the same row with different superscripts differ significantly ($P < 0.05$)

Table 4: Serum analysis of broiler chicks fed different levels of ATSM

Parameters	T1	T2	T3	T4	T5	SEM
MLA (U/mg Hb)	1.85 ^c	2.77 ^b	2.93 ^b	3.04 ^a	3.11 ^a	0.03
SDA (U/mg Hb)	35.7 ^b	39.8 ^b	40.7 ^a	43.5 ^a	45.3 ^a	1.21
GPx (U/mg Hb)	27.1 ^b	29.4 ^b	33.8 ^a	34.7 ^a	38.3 ^a	1.96
CAT (U/mg Hb)	54.2 ^a	45.7 ^b	42.5 ^b	41.6 ^b	40.1 ^b	0.52

Means in the same row with different superscripts differ significantly ($P < 0.05$)

Table 5: Antioxidant status of broiler chicks fed different levels of ATSM

Result and Discussion

The proximate composition of experimental diet (Table 1) revealed that it contains crude protein of 23.08 %, 20.11 % and 19.33 %; energy of 2936.0 kcal, 3000.8 kcal and 3100.2 kcal for starter, growers and finisher mash. The ether extract range between (4.28 – 5.03 %) and crude fibre range between (3.06 – 3.95 %). The proximate components meet the nutritional needs of birds according to NRC (1994). The crude fibre and ether extract range also conforms to the report of Teodora et al. (2020) in feeding broilers *Hermetia illucens* meal. The calcium (0.98 – 1.10 %) and phosphorus (0.47 – 0.51 %) range in the experimental diet is in line with the reports of Fascina et al. (2007); Aduku (2004). Proper feeding is one of the key cardinals of management in livestock production, for animals to perform at their optimum, there is need to furnish them with proper balanced diet which contains all the necessary nutrients (Alagbe and Oluwafemi, 2019).

The amino acid composition of *Albizia lebbek* stem bark is presented in Table 2. Results revealed the presence of threonine (1.12%), leucine (1.00 %), lysine (0.86 %), valine (0.65 %), glycine (1.00 %), phenylalanine (0.34 %), histidine (3.45 %), methionine (0.02 %), alanine (3.00 %), serine (0.78 %), proline (0.03 %), aspartic acid (2.00 %), arginine (1.65 %), tyrosine (0.10 %), isoleucine (2.00 %), aspartic acid (2.00 %) and cysteine (4.11 %). The sample contains high concentration of histidine and tyrosine has the lowest concentration. Amino acids are building blocks of protein which are necessary for gene expression and cell signal transduction regulation (Chzmruspollert et al., 2004). Phenylalanine plays a vital role in insulin secretion and fat oxidation (Ma et al., 2010). Lysine ensures effective production of hormones, enzymes and energy (Bazer et al., 2009). Alanine and glutamic acid enables a healthy skeletal system and energy production for the body Marc and Wu (2009); Kimura (2010). Regulation of blood sugar is been assisted by isoleucine (Tan et al., 2010; Yin et al., 2010). Adequate arginine ensures healthy immune system and maintains the visceral organs in the body (Brosnan and brosnan, 2010; Wu et al., 2010). Serine and cysteine play a key role as neuromodulator and antioxidant respectively (Wu et al., 2010; Wu et al., 2010; Baker, 2009). Methionine maintains the integrity of the liver, feather formation and egg size or production in birds (McKnight et al., 2010; Pali et al., 2009).

Haematological parameters of broiler chicks fed different levels of ATSM are presented in Table 4. PCV values ranged between (26.50 – 34.00 %), Hb (9.12 – 12.50 g/dl), RBC $1.88 - 2.97 (10^6/\mu\text{l})$, MCV (111.2 – 130.4 fl), MCH (34.01 – 59.00 pg) and MCHC (29.80 – 40.03 g/dl). RBC, PCV, Hb, MCV, MCH and MCHC values were higher ($P < 0.05$) in T3, T4 and T5 than for T2 and T1. WBC $20.41 - 30.04 (10^3/\mu\text{l})$ were highest in T4 and T5 ($P < 0.05$) compared to other treatments. Monocytes (0.07 – 1.72%), lymphocytes $10.45 - 20.04 (10^3/\mu\text{l})$, heterophils $1.23 - 6.11 (10^3/\mu\text{l})$ and eosinophils $0.88 - 1.27 (10^3/\mu\text{l})$ were lowest ($P < 0.05$) in T1 relative to other treatments. The haematological parameters measured follow similar pattern as it significantly ($P < 0.05$) increased from T1 to T5. However, all values are within the physiological range for normal birds (Talebi et al., 2005; Ibrahim, 2012; Subhadarsini and Silpa, 2020). Islam et al. (2004); Abdi-Hachesoo et al. (2011) reported a RBC range ($2.9 - 3.5 10^6/\mu\text{l}$), this variation could simply be as a result of differences

in age, sex, breed, environment, hormones and nutrition (Fudge, 2000). Haematological indices are used to in disease diagnosis as well as extent of damage to the blood (Nse Abasi et al., 2014; Omokore and Alagbe 2019). PCV and MCH are useful indices for the diagnosis of anaemia Nse Abasi et al. (2014); Alagbe (2019). A higher RBC level is an indication of adequate oxygen in the blood which gives room for effective nutrient transportation in the body (Isaac et al., 2013; Ugwuene, 2011). WBC plays a major role in the immune system by the production of antibodies, animals with low WBC stands a high risk of infection (Iwuji and Herbert, 2012; Isaac et al., 2013).

The serum biochemical indices of the experimental birds are presented in Table 5. Total protein ranges (2.57 – 3.97 g/dl), globulin (1.23 – 1.97 g/dl), albumin (1.34 – 2.03g/dl), creatinine (0.51 - 0.91 mg/dl) and glucose (196.1 – 250.6 mg/dl) were lowest ($P < 0.05$) for T1 compared to the other treatments while cholesterol (87.5 – 101.4 mg/dl), uric acid (50.7 – 74.1 mg/dl), ALT (50.7 – 74.1 u/l) and AST (200.9 – 300.7 u/l) was higher ($P < 0.05$) for T1 and T2 than for the rest of the treatments. Total protein value in T4 and T5 were significantly higher ($P < 0.05$) compared to the other treatment, this could be attributed to the presence of some relevant nutrients in ATSM. (Alagbe et al., 2020). Albumin content in the blood are generally influenced by protein shortages, however, the values reported fall within the range reported by (Subhadarsini and Silpa, 2020). Ibrahim (2012); Obajuluwa et al. (2020); Olafadehan et al. (2020); Livingston et al. (2020) reported a globulin and uric acid range of (1.6 – 1.9 g/dl) and (3.7-5.2 mg/dl) respectively. This result is also in agreement with the findings of Obikaonu et al. (2011) and Simaraks et al. (2004). Cholesterol, creatinine, uric acid, ALT and AST values follow similar pattern as it significantly ($P < 0.05$) decreased from T1 to T5. However, all the values were within the range reported by Olafadehan et al. (2020). Lower Creatinine and uric acid level is a sign that the kidney is not damage by feeding ATSM to the birds. According to Alagbe (2020), ATSM is loaded with several minerals, vitamins and bioactive chemicals or secondary metabolites (tannin, saponin, flavonoids, alkaloids, phenol etc.) which are within the lethal dose for broiler chicks. Urea levels is also reported to be influenced by dietary protein quality, quantity, bleeding time and are sensitive biomarkers employed in the diagnosis of renal damage (Akande and Odunsi, 2012). ATSM can also be serves as a hypolipidemic substance because of its ability to lower blood cholesterol, thus preventing heart diseases (Alagbe, 2020). ALT and AST are serum enzymes triggered due to the presence of a toxic substance in feed (Olabanji et al., 2007; Oluwafemi et al., 2020). The result obtain revealed that ATSM did not contain antinutrients or toxic substance which could hinder the general performance of birds, this result confirms the earlier report of Abdel et al. (2014); Cho et al. (2014) on the effects phytogetic feed additive in broiler chicks.

The oxidative status as influenced by ATSM is presented in Table 6. Superoxide dismutase [SDA; 35.7 – 45.3 U/mg Hb], glutathione peroxidase [GPx; 27.1 – 28.3 U/mg Hb], catalase [CAT; 40.1 – 54.2 U/mg Hb] and malonyldialdehyde [MLA; 1.85 – 3.11 U/mg Hb] values were lowest ($P < 0.05$) in T1 than in other treatments. According to Alagbe et al. (2019), ATSM contains antioxidants which are capable of scavenging free radicals, thereby giving total protection to animals. The presence of phenol and flavonoids prevent oxidative damage to biomolecules,

superoxide anions and lipid peroxy radicals (Hollman, 2001; Ojewuyi et al., 2014). The same results were reported by Lin et al. (2003) who observed that intake of phytogetic feed additives resulted in the increase in serum antioxidant enzyme activities and a decrease in MDA concentration. Conversely, Lan et al. (2013) reported that the concentration of blood glutathione was not affected by phytogetic feed additives.

Conclusion

Feed additives (plants extracts) have been reported to perform multiple biological activities including antibacterial, antifungal, antiviral, antihelminthic, antioxidant and immune modulator because they contain phytochemicals such as phenols, flavonoids, alkaloids, tannins, saponins, terpenoids etc. They are relatively cheap, safe and effective without any side effect on continuous use. The use of ATSM at 30 ml/litre of water have shown to be able to give total protection to the body and its metabolism against free radicals due to the presence of antioxidants and have no deleterious effect on the blood profile of broiler chicks.

References

- Teodora, P., Evgeni, P and Maya, I. (2020). Effect of black soldier fly (*Hermetia illucens*) meals on the meat quality of broilers. *Agriculture and Food Science*. 29:177-188.
- Aldi-Hachesoo, B., Talebi, A and Asri-Rezaei, S. (2012). Comparative study on blood profiles of indigenous and Ross-308 broiler breeders. *Global Veterinary Journal* 7:238-241.
- Cho, J.H., Kim, H.J and Kim, I.H (2014). Effects of phytogetic feed additive on growth performance, digestibility, blood metabolites, intestinal microbiota, meat colour and relative weight after oral challenge with *Clostridium perfringens* in broilers. *Livestock Science*, 160:82-88.
- Alagbe, J.O and Oluwafemi, R.A. (2019). Growth performance of weaner rabbits fed Noni (*Morinda citrifolia*) and *Moringa olifera* leaf mixture as partial replacement of soya bean meal. *International Journal of Advanced Biological and Biomedical Research*. 7(2): 185-195
- Abdel-Wareth, A.A., Lohakare, J.D. (2014). Effect of dietary supplementation of peppermint on performance, egg quality and serum metabolic profile of Hy-Line brown hens during the late laying period. *Animal Feed Science and Technology*, 197:114-120.
- Brosnan, J.T and Brosnan, M.E (2010). Creatine metabolism and the urea cycle. *Journal of Molecular Genetics and Metabolism*. 100:S49-52.
- Alagbe, J.O., Shittu, M.D and Eunice Abidemi Ojo (2020). Prospect of leaf extracts on the performance and blood profile of monogastric – A review. *International Journal of Integrated Education*. 3(7): 122-127.
- Gupta, M., Mazumder, U., Umar, T., Gomathi, P and Kumar, R (2004). Antioxidant and hepato-protective effects of *Bulhinaracemosa* against paracetamol and carbon tetrachloride induced liver damage in rats. *Iran Journal of Pharmacological Therapy* 3:12-20.
- Mc Donald, S.D., Prenzler, M and Robards, K (2001). Phenolic content and antioxidant activity of olive extracts. *Journal of Food Chemistry* 73:73-84.
- Sharma, Y., Bilbis, L.S., Lawal, M., Hassan, S.W., Abbas, A.Y.I and Isezuo, S.A (2007). Acute and sub chronic toxicity studies of crude aqueous extract of *Albizia chevalier*. *Asian Journal of Biochemistry* 2(4):224-236.
- Labaran, I., Lukman, O., Afolabi, A.A., Jamil, D and Umar, M (2016). Analysis of some phytochemical and minerals found in aqueous stem bark extract of *Albizia lebeck*. *Dutse Journal of Pure and Applied Sciences*. 2(1):231-237.
- Gary, D.B and Richard, D.M (2002). Interrelationship between nutrition and immunity. VM 139 Series at the Veterinary Medicine – Large Animal Clinical Sciences Dept, UF/IFAS Extension (2002).
- Hollman, P.C. (2001). Evidence for health benefits of plant phenols: Local or systemic effects. *Journal of Science Food Agriculture*, 81: 842-852.
- Alagbe, J.O. (2020). Performance, hematology and serum biochemical parameters of weaner rabbits fed different levels of fermented *Lagenaria breviflora* whole fruit extract. *Advances in Research and Reviews*, 2020, 1:5.
- Wu X, Ruan Z, Gao YL, Yin YL, Zhou XH, Wang L, Geng MM, Hou YQ, Wu G (2010). Dietary supplementation with L-arginine or N-carbamylglutamate enhances intestinal growth and heat shock protein-70 expression in weanling pigs fed a corn- and soybean meal-based diet. *Amino Acids*. 2010; 39:831–839.
- Ojewuyi, O.B, Ajiboye, T. O, Adebajo, E. O, Balogun, A, Mohammed, A.O (2014). Proximate composition, phytochemical and mineral contents of young and mature *Polyalthia longifolia* Sonn.leaves Fountain Journal of Natural and Applied Sciences: 2014; 3(1): 10 – 19.
- Baker D.H (2009). Advances in protein-amino acid nutrition of poultry. *Amino Acids*. 37:29–41.
- Wu G, Bazer FW, Burghardt RC, Johnson GA, Kim SW, Knabe DA, Li P, Li XL, McKnight JR, et al. (2010). Proline and hydroxyproline metabolism: implications for animal and human nutrition. *Amino Acids*. Epub 2010 Aug 10.
- Wu G, Bazer FW, Burghardt RC, Johnson GA, Kim SW, Knabe DA, Li XL, Satterfield MC, Smith SB, et al (2010). Functional amino acids in swine nutrition and production. In: Doppenberg J, editor. *Dynamics in animal nutrition*. Wageningen (The Netherlands): Wageningen Academic Publishers; p. 69–98.
- McKnight JR, Satterfield MC, Jobgen WS, Smith SB, Spencer TE, Meininger CJ, McNeal CJ, Wu G (2010). Beneficial effects of L-arginine on reducing obesity: potential mechanisms and important implications for human health. *Amino Acids*. 39:349–357.
- Tan B, Yin Y, Kong X, Li P, Li X, Gao H, Li X, Huang R, Wu G. (2010). L-Arginine stimulates proliferation and prevents endotoxin-induced death of intestinal cells. *Amino Acids*. 38:1227–1235.
- Yin YL, Yao K, Liu ZJ, Gong M, Ruan Z, Deng D, Tan BE, Liu ZQ, Wu G.(2010). Supplementing L-leucine to a low-protein diet increases tissue protein synthesis in weanling pigs. *Amino Acids*. Epub 2010 May 15.
- Palii S.S, Kays CE, Deval C, Bruhat A, Fafournoux P, Kilberg MS (2009). Specificity of amino acid regulated gene expression: analysis of gene subjected to either complete or single amino acid deprivation. *Amino Acids*. 37:79–88.
- Marc Rhoads J, Wu G (2009). Glutamine, arginine, and leucine signalling in the intestine. *Amino Acids*. 37:111–122.
- Kimura H (2010). Hydrogen sulphide: from brain to gut. *Antioxidant Redox Signal*. 12:1111–1123.
- Abdi-Hachesoo, B., Talebi, A and Asri-Rezaei, S. (2011). Comparative study on blood profile of indigenous and Ross-308 broiler breeders. *Global Veterinary Journal*, 7:238-241.
- Fugde, A.M. (2000). *Laboratory Medicine: Avian and Exotic Pets*. W.B Saunders Company, Philadelphia.
- Islam, M.S., Lucky, N.S., Literak, I., Ahad, A., Rahman, M.M and Siddiui, M.S.I. (2004). Changes of haematological parameters of fayoumi, assil and local chickens reared in Sylhet region in Bangladesh. *International Journal of Poultry Science*, 3:144-147.
- Obiakaonu, H.O., Okoli, I.C., Opara, M.N., Okoro, V.M.O., Ogbuewu, I.P., Etuk, E.B and Udedibie. A.B.I (2011).

- Haematological and serum biochemical indices of starter broilers fed neem leaf meal. *Online Journal of Animal and Feed Research*, 1(4):150-154.
30. Subhadarsini, M and Silpa, M.G. (2020). Comparative haematology and biochemical parameters of Indigenous broiler chicken. *International Journal of Scientific Technology Research*, 9(4):972-978.
 31. Santi, D.U and Kim, I.H (2017). Efficacy of phytogenic feed additive on performance production and health status of monogastric animals – A review. *Annals of Animal Science*. 17(4): 929-948.
 32. Talebi, A., Asri-Rezaei, S., Rozeh-Chai, R and Sahraei, R. (2005). Comparative studies on haematological values of broiler strains (Ross, Cobb, Arbo-acres and Arian). *International Journal of Poultry Science*, 4(8):573-579.
 33. Livingston, M.L., Cowieson, A.J., Crespo, R., Hoang, V., Nogal, B and Browning, M. (2020). Effect of broiler genetics, age and gender on performance and blood chemistry. *Heliyon* 6 (2020) e 04400.
 34. Omokore, E.O and Alagbe, J.O. (2019). Efficacy of dried *Phyllanthus amarus* leaf meal as an herbal feed additive on the growth performance, haematology and serum biochemistry of growing rabbits. *International Journal of Academic Research and Development*. 4(3): 97-104.
 35. Obajuluwa, O.V., Sanwo, K.A., Egbeyale, L.T and Fafiolu, O.A. (2020). Performance, blood profile and gut morphometry of broiler chickens fed diets supplemented with *Yohimbe* (*Pausynstalia yohimbe*) and *Larvacide*. *Journal of Veterinary and Animal Science*
 36. Mahipal, C., Ashak, K.P., Shalini, B., Narayan, D., Sunil, E.J and Kusumakar, S. (2015). Dietary supplementation of a novel phytogenic feed additive: effects on nutrient metabolism, antioxidant status and immune response of goats. *Animal Production Science*, 2:21-28.
 37. Olubukola, S.O., Anthony, J.A and Adewale, A (2015). Sub-chronic administration of methanolic whole fruit extracts of *Lagenaria breviflora* (Benth) induces mild toxicity in rats. *Journal of Pharmacognosy and natural product*. 11:516-521.
 38. Alagbe, J.O. (2019). Haematology, serum biochemistry, relative organ weight and bacteria count of broiler chicken given different levels of *Luffa aegyptiaca* leaf extracts. *International Journal of Advanced Biological and Biomedical Research*. 7(4):382-392.
 39. Nse Abasi, N.E., Mary, E.W., Uduak, A and Edem, E.A.O (2014). Haematological parameters and factors affecting their values. *Journal of Agricultural Science*, 2(1): 37-47.
 40. Isaac, L. J., Abah, G., Akpan, B and Ekaette, I.U (2013). Haematological properties of different breeds and sexes of rabbits. *Proceedings of the 18th Annual Conference of Animal Science Association of Nigeria*. (Pg. 24-27).
 41. Uwaya, J.O., Okunrobo, L.O and Igbinuwa, P.O (2016). *Albizia zygia*: A comparative investigation of phytochemical composition, proximate analysis and anti-seizure properties of methanol extracts and its leaves and stem bark. *Journal of Phytochemistry* 1(4): 23-27.
 42. Ugwuene, M.C (2011). Effect of dietary palm kernel meal for maize on the haematology and serum chemistry of broiler turkey. *Nigerian Journal of Animal Science* 13:93-103.
 43. Lin J, Hunkapiller AA, Layton AC, Chang Y.J, Robbins KR (2003) Response of intestinal microbiota to antibiotic growth promoters in chickens. *Foodborne Pathog Dis* 10:331–337.
 44. Lan, PTN, le Binh T, Benno Y (2013). Impact of two probiotic *Lactobacillus* strains feeding on fecal lactobacilli and weight gains in chicken. *J Gen Appl Microbiol* 49:29–36
 45. Oluwafemi, R.A., Isiaka Olawale and Alagbe, J.O. (2020). Recent trends in the utilization of medicinal plants as growth promoters in poultry nutrition- A review. *Research in: Agricultural and Veterinary Sciences*. 4(1): 5-11.
 46. Alagbe, J.O. (2019). Growth performance and haemato-biochemical parameters of broilers fed different levels of *Parkia biglobosa* leaf extracts. *Academic Journal of Life Sciences*. 5(12): 107 – 115.
 47. Abdel, J, Giallongo F, Frederick T, Pate J, Walusimbi S, Elias RJ, Wall EH, Bravo D, Hristov AN (2014) Effects of dietary *Capsicum oleoresin* on productivity and immune responses in lactating dairy cows. *J Dairy Sci* 98:6327–6339
 48. Cho, S, Alden N, Lee K (2014) .Pathways and functions of gut microbiota metabolism impacting host physiology. *Current Opinion on Biotechnology*. 36:137–145
 49. Olafadehan, O.A., Oluwafemi, R.A and Alagbe, J.O. (2020). Performance, haemato-biochemical parameters of broiler chicks administered *Rolfe* (*Daniellia oliveri*) leaf extract as an antibiotic alternative. *Advances in Research and Reviews*, 2020, 1:4.
 50. A.O.A.C.(2000). Association of Official Analytical Chemists. *Official Methods of Analysis 19th Edition* Washington, D.C Pages 69-77.
 51. Aduku, A.O (2004). Animal nutrition in the tropics: Feeds and feeding in monogastric and ruminant nutrition. *Journal of Applied Poultry Research*, 13: 628-638.
 52. Duncan, D.B. (1955). Multiple range and multiple F-test. *Biometrics* 11(1):1-42.
 53. National Research Council (1994). Nutrient requirement of poultry 9th Rev Edn, Washington D.C. National Academy Press.
 54. Akande, T and Odunsi, A.A (2012). Nutritive value and biochemical changes in broiler chickens fed detoxified castor kernel cake based diets. *African Journal of Biotechnology*, 11(12):2904-2911
 55. Olabanji, R.O., GO Farinu., JA Akinlade., OO Ojebiyi., AA Odunsi and AA Akingbade (2007). Studies on Haematological and Serum Biochemical Characteristics of Weaner Rabbits Fed Different Levels of Wild Sunflower (*Tithonia diversifolia* Hemsl A. Gray) Leaf- Blood Meal Mixture. *International Journal of Agriculture and Apiculture Research*. 4 (1&2): 80-89.
 56. Olafadehan, O.A., Oluwafemi, R.A and Alagbe, J.O (2020). Carcass quality, nutrient retention and caeca microbial population of broiler chicks administered *Rolfe* (*Daniellia oliveri*) leaf extract as an antibiotic alternative. *Journal of Drug Discovery*. 14(33):146-154.
 57. Karuppanan, K (2013). Phytopharmacological properties of *Albizia* species: A review. *International Journal of Pharmacy and Pharmacological Sciences* 10 (5): 3-9.
 58. Mohammed, F., Singh, P.P and Irchhaiya, R (2012). Review on *Albizia lebbbecka* potent herbal drugs. *International Research Journal of Pharmacy* 3(1): 63-68.
 59. Alagbe, J.O and Soares, D.M. (2018). Effects of feeding different levels of *Azolla pinnata*, *Polyalthia longifolia*, *Tithonia diversifolia*, *Moringa olifera*, *Azadiracta indica* leaf meal infusion as an organic supplement on the performance and nutrient retention of growing grass cutters. *Greener Journal of Agricultural Sciences*. 8(1):01-11.