

Otocephaly: Agnathia Microstomia Synotia Syndrome-A Case Report

Ahmed Mohammad Bakhit*, Mahmoud Kamal Moustafa, Mohammed Sayed Elyan, Easa Wali Umar Magfory
Neonatology department, Faculty of Medicine, Assiut University, Egypt

*Corresponding Author: Ahmed Mohammad Bakhit, Neonatology department, Faculty of Medicine, Assiut University, Egypt

Received date: July 21 2020; Accepted date: August 27, 2020; Published date: September 03, 2020

Citation: A M Bakhit, M K Moustafa, M S Elyan, E W U Magfory. (2020) Otocephaly: Agnathia Microstomia Synotia Syndrome-A Case Report. International Journal of Clinical Case Reports and Reviews. 3(2); DOI: 10.31579/2690-4861/035

Copyright: © 2020 Ahmed Mohammad Bakhit, This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Abstract

Otocephaly is characterized by aplasia or hypoplasia of mandible, a small oral fissure and low lying ears at the level of neck usually meeting in the midline. This is due to the failure of migration of the ventral part of the first brachial arch. This rare lethal non familial condition occurs in 1 in 70,000 births. This is the first reported case of otocephaly in Tema city, Sohag governorate, Egypt as per our best knowledge, a baby girl born to non-consanguineous parents, who was unable to survive a long time after birth. This case is being reported because of its rarity.

Keywords: agnathia; antenatal diagnosis; otocephaly

Case Report

A 40yrs old female patient p1+0, 1 living male, 2ndry infertility 10 years, Previous failed ICSI, pregnant +_ 34 weeks, by ICSI (2nd trial). She admitted to a private centre for Women, Tema, Sohag, Egypt due to severe abdominal pain. There is no consanguinity, there was no exposure to teratogens or infection during the pregnancy. Dating scan was done and cervical cerclage was put at 14th week of gestation. Polyhydramnios was noted on ultrasound scan at 24th week of gestation.

Detail ultrasound scan on this admission revealed a single live fetus with an amniotic fluid index of 43 cm and an estimated fetal weight of 2250 g. There

was no definite fetal stomach noted which led to a suspicion of a possibility of esophageal atresia. Baby girl was born by cesarian section with birth weight of 2500 g. Examination of the baby revealed downwards slanted eyes, hypertelorism, very small oral aperture (Microstomia) with blind ended pouch confirmed by inability to pass the orogastric tube, absence of mandible (Agnathia) and external ears displaced ventromedially in the neck close to the mid line (Synotia) (**Figure 1**). Rest of the clinical examination was normal. Baby died soon after the birth without any active resuscitation, considering the gross abnormalities. Autopsy was advised but refused by the parents. Mother was discharged on following day after counselling with family planning advices.



Figure 1: Shows microstomia, absence of mandible (agnathia) and low-set ventromedially placed ears.

Discussion

Otocephaly is a rare lethal syndrome of microstomia, agnathia and ear anomalies. Other anomalies associated are holoprosencephaly, cephalocele, dysgenesis of corpus colossus, atresia of third ventricle, skeletal, genitourinary, cardiovascular system and endocrine gland hypoplasia [1, 2]. The incidence of otocephaly which is a rare idiopathic malformation in humans is 1 per 70,000 new-borns [3]. Every pharyngeal arch has a core of cartilage, unsegmented mesoderm, an artery from aortic arch and a nerve that enters it from the brain stem carrying motor fibers to supply the skeletal muscle derived from the mesoderm [1]. Arrest in the development of the first branchial arch due to an insult to the neural crest cells has been proposed to be the cause of this condition, which may be induced by chromosomal mutation or teratogens. Otocephaly, with associated anomalies are considered lethal due to severe respiratory dysfunction. Otocephaly is usually suspected on radiological antenatal checkup specially during the third trimester when it is impossible to visualize the mandible and ears are in a very low and in medial position [2,4]. Polyhydramnios may be the presenting feature during pregnancy [3]. In Sri Lanka the termination of the pregnancy is only considered if the life of the mother is in danger due to the pregnancy, so the antenatal identification of such cases over helps the parents to prepare for the delivery. The syndrome complex of otocephaly is divided into four types:

1. Isolated agnathia;
2. Agnathia with holoprosencephaly;
3. Agnathia with situs inversus and visceral anomalies;
4. Agnathia, holoprosencephaly, situs inversus and other visceral anomalies [5].

The case we describe belongs to type 2. Otocephaly anomaly shows spectrum of various manifestations ranges in severity from severe micrognathia as a part of the robin sequence to cyclopi holoprosencephaly complex invariably associated with fetal death [6]. There are few proposed teratogenic effects as the cause for

otocephaly such as exposure streptogramin antibiotics and trypan blue, theophylline [7]. In our case there was no such exposure identified. The genetic basis of otocephaly is still largely unclear. Even though otocephaly is a lethal syndrome there are cases on successful management of otocephalic babies after birth. Enteral feeding was successful in such newborns with isolated agnathia with nasogastric tubes [6].

Conclusion

Prenatal diagnosis of this lethal condition is possible, and it should be considered as one of the differential diagnosis for polyhydramnios. Poor prognosis of this condition should be discussed with parents prior the delivery. There is still no solid evidence to the cause of this rare condition.

References

1. Sharma M, Arora AK, Kohli D, Kohli S (2013) Otocephaly - A case report. *J Asian Med Students Assoc*.
2. Jagtap SV, Saini N, Jagtap S, Saini S (2015) Otocephaly: Agnathia-Microstomia-synotia syndrome-A rare congenital anomaly. *J Clin Diagnostic Res* 9: 3-4.
3. Faye-Petersen O, David E, Rangwala N, Seaman JP, Hua Z, et al. (2006) Otocephaly: Report of five new cases and a literature review. *Fetal Pediatr Pathol* 25: 277-296.
4. Ducarme G, Largilliere C, Amarenco B (2007) Three-dimensional ultrasound in prenatal diagnosis of isolated otocephaly. *Prenat Diagn* 27: 481-483.
5. Croen LA, Shaw GM, Lammer EJ (1996) Holoprosencephaly: Epidemiologic and clinical characteristics of a California population. *Am J Med Genet* 64: 465-472.
6. Sharma M, Case OA, Sharma A, Amrutha KV, Abraham J (2016) International journal of otocephaly: A case report. *Int J Res Med Sci* 4: 1286-1289.
7. Ibba RM, Zoppi MA, Floris M (2000) Otocephaly: Prenatal diagnosis of a new case and etiopathogenetic considerations. *Am J Med Genet* 90: 427-429.



This work is licensed under Creative Commons Attribution 4.0 License

To Submit Your Article Click Here: [Submit Article](#)

DOI: [10.31579/2690-4861/035](https://doi.org/10.31579/2690-4861/035)

Ready to submit your research? Choose Auctores and benefit from:

- ❖ fast, convenient online submission
- ❖ rigorous peer review by experienced research in your field
- ❖ rapid publication on acceptance
- ❖ authors retain copyrights
- ❖ unique DOI for all articles
- ❖ immediate, unrestricted online access

At Auctores, research is always in progress.

Learn more www.auctoresonline.org/journals/international-journal-of-clinical-case-reports-and-reviews