

An Exaggerated Capillary Endothelial Edema May be the Cause of Sudden Deaths in Sickle Cell Diseases

Mehmet Rami Helvaci¹, Kubra Piral², Kubra Seckin³, Esin Esra Tanaydin³, Ayse Deniz Karabacak³, Mehpare Camlibel⁴, Abdulrazak Abyad⁵, Lesley Pocock⁶

¹ Specialist of Internal Medicine, MD, Turkey.

² Manager of Writing and Statistics, Turkey.

³ Ministry of Health of Turkey, MD, Turkey.

⁴ Specialist of Emergency Medicine, MD, Turkey.

⁵ Middle-East Academy for Medicine of Aging, MD, Lebanon.

⁶ Medi-WORLD International, Australia.

***Corresponding Author:** Mehmet Rami Helvaci, Specialist of Internal Medicine, MD, Turkey.

Received date: June 01, 2023; **Accepted date:** June 09, 2023; **Published date:** June 21, 2023

Citation: Mehmet R. Helvaci, Kubra Piral, Kubra Seckin, Esin Esra Tanaydin, Ayse Deniz Karabacak, et al, (2023), An Exaggerated Capillary Endothelial Edema May be the Cause of Sudden Deaths in Sickle Cell Diseases. *J. Gastroenterology Pancreatology and Hepatobiliary Disorders*, 7(1); DOI:10.31579/2641-5194/065

Copyright: © 2023, Mehmet Rami Helvaci. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract

Background: Sickle cell diseases (SCDs) are inborn and severe inflammatory processes on vascular endothelium, particularly at the capillaries which are the actual distributors of the sickled or just hardened red blood cells (RBCs) into the tissues.

Methods: All patients of the SCDs were included.

Results: We studied 222 males and 212 females with similar ages (30.8 vs 30.3 years, $p>0.05$, respectively). Disseminated teeth losses (5.4% vs 1.4%, $p<0.001$), ileus (7.2% vs 1.4%, $p<0.001$), cirrhosis (8.1% vs 1.8%, $p<0.001$), leg ulcers (19.8% vs 7.0%, $p<0.001$), digital clubbing (14.8% vs 6.6%, $p<0.001$), coronary heart disease (18.0% vs 13.2%, $p<0.05$), chronic renal disease (9.9% vs 6.1%, $p<0.05$), chronic obstructive pulmonary disease (25.2% vs 7.0%, $p<0.001$), and stroke (12.1% vs 7.5%, $p<0.05$) were higher in males but not acute chest syndrome (2.7% vs 3.7%), pulmonary hypertension (12.6% vs 11.7%), deep venous thrombosis and/or varices and/or telangiectasias (9.0% vs 6.6%), or mean age of mortality (30.2 vs 33.3 years) ($p>0.05$ for all).

Conclusion: The sickled or just hardened RBCs-induced capillary endothelial damage, inflammation, edema, and fibrosis are initiated at birth, and terminate with disseminated tissue hypoxia, multiorgan failures, and sudden deaths even at childhood. Although RBCs suspensions and corticosteroids in acute and aspirin plus hydroxyurea in acute and chronic phases decrease severity of the destructive process, survivals are still shortened in both genders, dramatically. Infections, medical or surgical emergencies, or emotional stresses-induced increased basal metabolic rate aggravates the sickling and capillary endothelial edema, and may terminate with multiorgan failures-induced sudden deaths in the SCDs.

Key words: sickle cell diseases; sickled or just hardened red blood cells; capillary endothelial damage; exaggerated capillary endothelial edema; sudden deaths

Introduction

Chronic endothelial damage may be the main underlying cause of aging and death by causing end-organ failures [1]. Much higher blood pressures (BPs) of the afferent vasculature may be the chief accelerating factor by causing recurrent injuries on vascular endothelium. Probably, whole

afferent vasculature including capillaries are mainly involved in the destructive process. Thus the term of venosclerosis is not as famous as atherosclerosis in the literature. Due to the chronic endothelial damage, inflammation, edema, and fibrosis, vascular walls thicken, their lumens

narrow, and they lose their elastic natures which eventually reduce blood flow to the terminal organs, and increase systolic and decrease diastolic BPs further. Some of the well-known accelerating factors of the harmful process are physical inactivity, sedentary lifestyle, animal-rich diet, smoking, alcohol, overweight, chronic inflammations, prolonged infections, and cancers for the development of terminal consequences including obesity, hypertension (HT), diabetes mellitus (DM), cirrhosis, chronic obstructive pulmonary disease (COPD), coronary heart disease (CHD), chronic renal disease (CRD), stroke, peripheral artery disease (PAD), mesenteric ischemia, osteoporosis, dementia, early aging, and premature death [2, 3]. Although early withdrawal of the accelerating factors can delay terminal consequences, after development of obesity, HT, DM, cirrhosis, COPD, CRD, CHD, stroke, PAD, mesenteric ischemia, osteoporosis, aging, and dementia-like end-organ insufficiencies, the endothelial changes can not be reversed due to their fibrotic natures, completely. The accelerating factors and terminal consequences of the harmful process are researched under the titles of metabolic syndrome, aging syndrome, and accelerated endothelial damage syndrome in the literature [4-6]. On the other hand, sickle cell diseases (SCDs) are inborn and severe inflammatory and highly destructive processes on vascular endothelium, initiated at birth and terminated with an advanced atherosclerosis-induced end-organ failures in much earlier ages of life [7, 8]. Hemoglobin S causes loss of elastic and biconcave disc shaped structures of red blood cells (RBCs). Probably loss of elasticity instead of shape is the major problem because sickling is rare in peripheral blood samples of the patients with associated thalassemia minors (TMs), and human survival is not affected in hereditary spherocytosis or elliptocytosis. Loss of elasticity is present even at birth, but exaggerated with inflammations, infections, and emotional stresses of the body. The sickled or just hardened RBCs-induced chronic endothelial damage, inflammation, edema, and fibrosis terminate with disseminated tissue hypoxia all over the body [9]. As a difference from other causes of chronic endothelial damage, SCDs keep vascular endothelium particularly at the capillaries which are the actual distributors of the sickled or just hardened RBCs into the tissues [10, 11]. The sickled or just hardened RBCs-induced chronic endothelial damage builds up an advanced atherosclerosis in much earlier ages of life. Vascular narrowings and occlusions-induced tissue ischemia and end-organ failures are the terminal results, so the life expectancy is decreased by 25 to 30 years for both genders in the SCDs [8].

Material and methods

The clinical study was performed in Medical Faculty of the Mustafa Kemal University between March 2007 and June 2016. All patients of the SCDs were included. The SCDs are diagnosed with the hemoglobin electrophoresis performed by means of high performance liquid chromatography (HPLC). Smoking and alcohol habits, acute painful crises per year, transfused units of RBCs in their lives, leg ulcers, stroke, surgical operations, deep venous thrombosis (DVT), epilepsy, and priapism were learnt. Cases with a history of one pack-year were accepted as smokers, and one drink-year were accepted as drinkers. A complete physical examination was performed by the Same Internist, and patients with disseminated teeth losses (<20 teeth present) were detected. Patients with an acute painful crisis or any other inflammatory process were treated at first, and the laboratory tests and clinical measurements were performed on the silent phase. A check up procedure including serum

iron, iron binding capacity, ferritin, creatinine, liver function tests, markers of hepatitis viruses A, B, and C, marker of human immunodeficiency virus, a posterior-anterior chest x-ray film, an electrocardiogram, a Doppler echocardiogram both to evaluate cardiac walls and valves, and to measure systolic BPs of pulmonary artery, an abdominal ultrasonography, a venous Doppler ultrasonography of the lower limbs, a computed tomography (CT) of brain, and a magnetic resonance imaging (MRI) of hips was performed. Other bones for avascular necrosis were scanned according to the patients' complaints. So avascular necrosis of bones was diagnosed via MRI [12]. Associated TMs were detected with serum iron, iron binding capacity, ferritin, and hemoglobin electrophoresis performed via HPLC, because the SCDs with associated TMs show a milder clinic than the sickle cell anemia (SCA) (Hb SS) alone [13]. Systolic BPs of the pulmonary artery of 40 mmHg or higher are accepted as pulmonary hypertension (PHT) [14]. The criterion for diagnosis of COPD is a post-bronchodilator forced expiratory volume in one second/forced vital capacity of lower than 70% [15]. Acute chest syndrome (ACS) is diagnosed clinically with the presence of new infiltrates on chest x-ray film, fever, cough, sputum, dyspnea, or hypoxia [16]. An x-ray film of abdomen in upright position was taken in patients with abdominal distention or discomfort, vomiting, obstipation, or lack of bowel movement, and ileus is diagnosed with gaseous distention of isolated segments of bowel, vomiting, obstipation, cramps, and with the absence of peristaltic activity. CRD is diagnosed with a persistent serum creatinine level of 1.3 mg/dL or greater in males and 1.2 mg/dL or greater in females. Cirrhosis is diagnosed with physical examination, laboratory parameters, and ultrasonographic findings. Clubbing is diagnosed with the ratio of distal phalangeal diameter to interphalangeal diameter of greater than 1.0, and with the presence of Schamroth's sign [17, 18]. An exercise electrocardiogram is performed in cases with an abnormal electrocardiogram and/or angina pectoris. Coronary angiography is performed for the cases with exercise electrocardiogram positivity. So CHD is diagnosed either angiographically or with the Doppler echocardiographic findings as movement disorders in the cardiac walls. Rheumatic heart disease is diagnosed with the echocardiographic findings, too. Stroke is diagnosed by the CT. Sickle cell retinopathy is diagnosed with ophthalmologic examination in cases with visual complaints. Mann-Whitney U test, Independent-Samples t test, and comparison of proportions were the methods of statistical analyses.

Results

The study included 222 males and 212 females with similar ages (30.8 vs 30.3 years, $p>0.05$, respectively). Prevalences of associated TMs were similar in both genders, too (72.5% vs 67.9%, $p>0.05$, respectively). Smoking (23.8% vs 6.1%) and alcohol (4.9% vs 0.4%) were higher in males ($p<0.001$ for both) (**Table 1**). Transfused units of RBCs in their lives (48.1 vs 28.5, $p=0.000$), disseminated teeth losses (5.4% vs 1.4%, $p<0.001$), ileus (7.2% vs 1.4%, $p<0.001$), cirrhosis (8.1% vs 1.8%, $p<0.001$), leg ulcers (19.8% vs 7.0%, $p<0.001$), clubbing (14.8% vs 6.6%, $p<0.001$), CHD (18.0% vs 13.2%, $p<0.05$), CRD (9.9% vs 6.1%, $p<0.05$), COPD (25.2% vs 7.0%, $p<0.001$), and stroke (12.1% vs 7.5%, $p<0.05$) were higher in males but not ACS (2.7% vs 3.7%), PHT (12.6% vs 11.7%), DVT and/or varices and/or telangiectasias (9.0% vs 6.6%), or mean age of mortality (30.2 vs 33.3 years) ($p>0.05$ for all) (**Table 2**). Beside that the mean ages of terminal consequences were shown in **Table 3**.

Variables	Males with SCDs*	p-value	Females with SCDs
Prevalence	51.1% (222)	Ns†	48.8% (212)
Mean age (year)	30.8 ± 10.0 (5-58)	Ns	30.3 ± 9.9 (8-59)
Associated TMs‡	72.5% (161)	Ns	67.9% (144)
Smoking	23.8% (53)	<0.001	6.1% (13)
Alcoholism	4.9% (11)	≤0.001	0.4% (1)

Table 1: Characteristic features of the study patients

*Sickle cell diseases †Nonsignificant (p>0.05) ‡Thalassemia minor

Variables	Males with SCDs*	p-value	Females with SCDs
Painful crises per year	5.0 ± 7.1 (0-36)	Ns†	4.9 ± 8.6 (0-52)
Transfused units of RBCs‡	48.1 ± 61.8 (0-434)	0.000	28.5 ± 35.8 (0-206)
Disseminated teeth losses (<20 teeth present)	5.4% (12)	<0.001	1.4% (3)
COPD§	25.2% (56)	<0.001	7.0% (15)
Ileus	7.2% (16)	<0.001	1.4% (3)
Cirrhosis	8.1% (18)	<0.001	1.8% (4)
Leg ulcers	19.8% (44)	<0.001	7.0% (15)
Digital clubbing	14.8% (33)	<0.001	6.6% (14)
CHD¶	18.0% (40)	<0.05	13.2% (28)
CRD**	9.9% (22)	<0.05	6.1% (13)
Stroke	12.1% (27)	<0.05	7.5% (16)
PHT***	12.6% (28)	Ns	11.7% (25)
Autosplenectomy	50.4% (112)	Ns	53.3% (113)
DVT**** and/or varices and/or telangiectasias	9.0% (20)	Ns	6.6% (14)
Rheumatic heart disease	6.7% (15)	Ns	5.6% (12)
Avascular necrosis of bones	24.3% (54)	Ns	25.4% (54)
Sickle cell retinopathy	0.9% (2)	Ns	0.9% (2)
Epilepsy	2.7% (6)	Ns	2.3% (5)
ACS*****	2.7% (6)	Ns	3.7% (8)
Mortality	7.6% (17)	Ns	6.6% (14)
Mean age of mortality (year)	30.2 ± 8.4 (19-50)	Ns	33.3 ± 9.2 (19-47)

Table 2: Associated pathologies of the study patients

*Sickle cell diseases †Nonsignificant (p>0.05) ‡Red blood cells §Chronic obstructive pulmonary disease ¶Coronary heart disease **Chronic renal disease ***Pulmonary hypertension ****Deep venous thrombosis *****Acute chest syndrome

Variables	Mean age (year)
Ileus	29.8 ± 9.8 (18-53)
Hepatomegaly	30.2 ± 9.5 (5-59)
ACS*	30.3 ± 10.0 (5-59)
Sickle cell retinopathy	31.5 ± 10.8 (21-46)
Rheumatic heart disease	31.9 ± 8.4 (20-49)
Autosplenectomy	32.5 ± 9.5 (15-59)
Disseminated teeth losses (<20 teeth present)	32.6 ± 12.7 (11-58)
Avascular necrosis of bones	32.8 ± 9.8 (13-58)
Epilepsy	33.2 ± 11.6 (18-54)
Priapism	33.4 ± 7.9 (18-51)
Left lobe hypertrophy of the liver	33.4 ± 10.7 (19-56)
Stroke	33.5 ± 11.9 (9-58)
COPD†	33.6 ± 9.2 (13-58)
PHT‡	34.0 ± 10.0 (18-56)
Leg ulcers	35.3 ± 8.8 (17-58)
Digital clubbing	35.4 ± 10.7 (18-56)
CHD§	35.7 ± 10.8 (17-59)
DVT¶ and/or varices and/or telangiectasias	37.0 ± 8.4 (17-50)
Cirrhosis	37.0 ± 11.5 (19-56)
CRD**	39.4 ± 9.7 (19-59)

Table 3: Mean ages of the consequences of the sickle cell diseases

*Acute chest syndrome †Chronic obstructive pulmonary disease ‡Pulmonary hypertension §Coronary heart disease ¶Deep venous thrombosis **Chronic renal disease

Discussion

Acute painful crises are the most disabling symptoms of the SCDs. Although some authors reported that pain itself may not be life threatening directly, infections, medical or surgical emergencies, or

emotional stresses are the most common precipitating factors of the crises [19]. Although the sickled or just hardened RBCs-induced capillary endothelial damage, inflammation, and edema are present even at birth, the increased basal metabolic rate during such stresses aggravates the sickling and capillary endothelial damage, inflammation, and edema, and

may terminate with disseminated tissue hypoxia and multiorgan failures-induced sudden deaths in the SCDs. So the risk of mortality is much higher during the crises. Actually, each crisis may complicate with the following crises by leaving some sequelae on the capillary endothelial system all over the body. After a period of time, the sequelae may terminate with sudden end-organ failures and death during a final acute painful crisis that may even be silent, clinically. Similarly, after a 20-year experience on such patients, the deaths seem sudden and unexpected events in the SCDs. Unfortunately, most of the deaths develop just after the hospital admission, and majority of such cases are without hydroxyurea therapy [20]. Rapid RBCs supports are usually life-saving for such patients, although preparation of RBCs units for transfusion usually takes time. Beside that RBCs supports in emergencies become much more difficult in such terminal patients due to the repeated transfusions-induced blood group mismatch. Actually, transfusion of each unit of RBCs complicates the following transfusions by means of the blood subgroup mismatch. Due to the significant efficacy of hydroxyurea therapy, RBCs transfusions should be kept just for acute events and emergencies in the SCDs [21]. According to our experiences, simple and repeated transfusions are superior to RBCs exchange in the SCDs [22]. First of all, preparation of one or two units of RBCs suspensions in each time rather than preparation of six units or higher provides time to clinicians to prepare more units by preventing sudden death of such high-risk cases. Secondly, transfusions of one or two units of RBCs suspensions in each time decrease the severity of pain and relax anxiety of the patients and their relatives because RBCs transfusions probably have the strongest analgesic effects during such crises. Actually, the decreased severity of pain by transfusions also indicates the decreased severity of inflammation in whole body. Thirdly, transfusions of lesser units of RBCs suspensions in each time by means of the simple transfusions decrease transfusions-related complications including infections, iron overload, and blood group mismatch. Fourthly, transfusions of RBCs suspensions in the secondary health centers prevent some deaths developed during the transport to the tertiary centers for the exchange. Finally, cost of the simple and repeated transfusions on insurance system is much lower than the exchange that needs trained staff and additional devices. On the other hand, pain is the result of complex and poorly understood interactions between RBCs, white blood cells (WBCs), platelets (PLTs), and endothelial cells, yet. Whether leukocytosis contributes to the pathogenesis by releasing cytotoxic enzymes is unknown. The adverse actions of WBCs on the capillary endothelium are of particular interest with regard to the cerebrovascular diseases in the SCDs. For instance, leukocytosis even in the absence of an infection was an independent predictor of the severity of the SCDs, and it was associated with the higher risk of stroke [23]. Disseminated tissue hypoxia, releasing of inflammatory mediators, bone infarctions, and activation of afferent nerves may take role in the pathophysiology of the intolerable pain. Because of the severity of pain, narcotic analgesics are usually required to control them [24], but according to our long term experience, simple and repeated RBCs transfusions are much more effective than the narcotics to control the intolerable pain in the SCDs.

Hydroxyurea is the first drug that was approved by Food and Drug Administration in the SCDs [25]. It is an orally-administered, cheap, safe, and effective drug, and it may be the only life-saving drug in the treatment of the SCDs [26, 27]. It interferes with the cell division by blocking the formation of deoxyribonucleotides via inhibition of ribonucleotide reductase. The deoxyribonucleotides are the building blocks of DNA. Hydroxyurea mainly affects hyperproliferating cells. Although the action way of hydroxyurea is thought to be the increase in gamma-globin synthesis for fetal hemoglobin (Hb F), its main action may be the prevention of leukocytosis and thrombocytosis by blocking the DNA synthesis [28, 29]. By this way, the inborn inflammatory and destructive process of the SCDs is suppressed with some extent. Due to the same action way, hydroxyurea is also used in moderate and severe psoriasis to suppress hyperproliferating skin cells. As also seen in the viral hepatitis

cases, although presence of a continuous damage of sickled or just hardened RBCs on the capillary endothelium, the severity of destructive process may be exaggerated by the patients' own WBCs and PLTs. So suppression of proliferation of the WBCs and PLTs may limit the capillary endothelial damage, inflammation, edema, tissue ischemia, and end-organ failures in the body [30]. Similarly, final Hb F levels in the hydroxyurea users did not differ from their pretreatment levels [31]. The Multicenter Study of Hydroxyurea (MSH) studied 299 severely affected adults with the SCA, and compared the results of patients treated with hydroxyurea or placebo [32]. The study particularly researched effects of hydroxyurea on the painful crises, ACS, and requirement of RBCs transfusion. The outcomes were so overwhelming in the favour of hydroxyurea that the study was terminated after 22 months, and hydroxyurea was started for all patients. The MSH also demonstrated that patients treated with hydroxyurea had a 44% decrease in hospitalizations [32]. In multivariable analyses, there was a strong and independent association of lower neutrophil counts with the lower crisis rates [32]. But this study was performed just in severe SCA cases alone, and the rate of painful crises was decreased from 4.5 to 2.5 per year [32]. Whereas we used all subtypes of the SCDs with all clinical severity, and the rate of painful crises was decreased from 10.3 to 1.7 per year ($p < 0.000$) with an additional decreased severity of them (7.8/10 vs 2.2/10, $p < 0.000$) (27). Parallel to our results, adults using hydroxyurea therapy for frequent painful crises appear to have a reduced mortality rate after a 9-year follow-up period [33]. Although the underlying disease severity remains critical to determine prognosis, hydroxyurea may also decrease severity of the SCDs and prolong survival [33]. The complications start to be seen even in infancy in the SCDs. For instance, infants with lower hemoglobin values were more likely to have higher incidences of clinical events such as ACS, acute painful crises, and lower neuropsychological scores, and hydroxyurea reduced the incidences of them [34]. Hydroxyurea therapy in early years of life may improve growth, and prevent end-organ failures. Transfusion programmes can also reduce all of the complications, but transfusions carry many risks including infections, iron overload, and development of allo-antibodies causing subsequent transfusions difficult. On the other hand, elevations of liver enzymes during some acute painful crises can not be reversed by withdrawing of the hydroxyurea therapy alone, instead withdrawal of all of the medications were highly effective in such cases during the 20-year experience on such patients. After normalization of the liver enzymes, the essential medications must be started one by one, instead of all of them at the same time, again. Thus hydroxyurea must even be used during the acute painful crises. Additionally, we observed mild, moderate, or even severe bone marrow suppressions and pancytopenia in some patients using high-dose hydroxyurea (35 mg/kg/day). Interestingly, such cases were completely silent other than some signs and symptoms of anemia, and all of them were resolved completely just by giving a few-day break for the hydroxyurea therapy and starting with smaller doses again.

Aspirin is a nonsteroidal anti-inflammatory drug (NSAID) used to reduce inflammation and acute thromboembolic events. Although aspirin has similar anti-inflammatory effects with the other NSAIDs, it also suppresses the normal functions of PLTs, irreversibly. This property causes aspirin being different from other NSAIDs, which are reversible inhibitors. Aspirin acts as an acetylating agent where an acetyl group is covalently attached to a serine residue in the active site of the cyclooxygenase (COX) enzyme. Aspirin's ability to suppress the production of prostaglandins (PGs) and thromboxanes (TXs) is due to its irreversible inactivation of the COX enzyme required for PGs and TXs synthesis. PGs are the locally produced hormones with some diverse effects, including the transmission of pain into the brain and modulation of the hypothalamic thermostat and inflammation. TXs are responsible for the aggregation of PLTs to form blood clots. In another definition, low-dose aspirin use irreversibly blocks the formation of TXA₂ in the PLTs, producing an inhibitory effect on the PLT aggregation during whole lifespan of the affected PLTs (8-9 days). Since PLTs do not have

nucleus and DNA, they are unable to synthesize new COX enzyme once aspirin inhibited the enzyme. The antithrombotic property of aspirin is useful to reduce the incidences of myocardial infarction, transient ischemic attack, and stroke [35]. Heart attacks are caused primarily by blood clots, and low dose of aspirin is seen as an effective medical intervention to prevent a second myocardial infarction [36]. According to the medical literature, aspirin may also be effective in prevention of colorectal cancers [37]. On the other hand, aspirin has some side effects including gastric ulcers, gastric bleeding, worsening of asthma, and Reye syndrome in childhood and adolescence. Reye syndrome is a rapidly worsening brain disease [38]. The first detailed description of Reye syndrome was in 1963 by an Australian pathologist, Douglas Reye [39]. The syndrome mostly affects children, but it can only affect fewer than one in a million children a year [39]. It usually starts just after recovery from a viral infection, such as influenza or chicken pox [39]. Symptoms of Reye syndrome may include personality changes, confusion, seizures, and loss of consciousness (38). Although the liver toxicity typically occurs in the syndrome and the liver is enlarged in most cases, jaundice is usually not seen with it [38]. Early diagnosis improves outcomes, and treatment is supportive. Mannitol may be used in cases with the brain swelling [39]. Although the death occurs in 20-40% of patients, about one third of survivors get a significant degree of brain damage (38). Interestingly, about 90% of cases in children are associated with an aspirin use [40]. Due to the risk of Reye syndrome, the US Food and Drug Administration recommends that aspirin or aspirin-containing products should not be prescribed for febrile patients under the age of 16 years [41]. Eventually, the general recommendation to use aspirin in children has been withdrawn, and it was only recommended for Kawasaki disease [38]. When aspirin use was withdrawn for children in the US and UK in the 1980s, a decrease of more than 90% of Reye syndrome was seen [39]. Due to the higher side effects of corticosteroids in long term, and due to the very low risk of Reye syndrome but much higher risk of death due to the SCDs even in children, aspirin should be added into the acute and chronic phase treatments of the SCDs with an anti-inflammatory dose even in childhood [42].

ACS is a significant cause of mortality in the SCDs [43]. It occurs most often as a single episode, and a past history is associated with a higher mortality rate [43]. Similarly, all of 14 patients with ACS had just a single episode, and two of them were fatal in spite of the immediate RBCs and ventilation supports and antibiotic therapy in the present study. The remaining 12 patients are still alive without a recurrence at the end of the 10-year follow up period. ACS is the most common between two to four years of age, and its incidence decreases with aging [44]. As a difference from atherosclerotic consequences, the incidence of ACS did not show an increase with aging in the present study, and the mean ages of the patients with ACS and SCDs were similar (30.3 vs 30.5 years, $p>0.05$, respectively). The decreased incidence with aging may be due to the high mortality rate during the first episode and/or an acquired immunity against various antigens, and/or decreased strength of immune response by aging. Probably, ACS shows an inborn severity of the SCDs, and the incidence of ACS is higher in severe patients such as patients with the SCA and higher WBCs counts [43, 44]. According to our long term experiences on the SCDs, the increased metabolic rate during infections accelerates sickling, thrombocytosis, leukocytosis, and capillary endothelial damage and edema, and terminates with end-organ failures-induced sudden deaths. ACS may also be a collapse of the pulmonary vasculature during such infections, and the exaggerated immune response against the sickled or just hardened RBCs-induced diffuse capillary endothelial damage may be important in the high mortality rate. A preliminary result from the Multi-Institutional Study of Hydroxyurea in the SCDs indicating a significant reduction of episodes of ACS with hydroxyurea therapy suggests that a considerable number of episodes are exaggerated with the increased numbers of WBCs and PLTs (45). Similarly, we strongly recommend hydroxyurea for all patients that may also be the cause of low incidence of ACS in our follow up cases (2.7%

in males and 3.7% in females). Additionally, ACS did not show an infectious etiology in 66% (43, 44), and 12 of 27 cases with ACS had evidence of fat embolism in the other study (46). Beside that some authors indicated that antibiotics did not shorten the clinical course (47). RBCs support must be given as earliest as possible. RBCs support has the obvious benefits of decreasing sickle cell concentration directly, and suppressing bone marrow for the production of abnormal RBCs and excessive WBCs and PLTs. So they prevent further sickling-induced exaggerated capillary endothelial edema, disseminated tissue hypoxia, and end-organ failures-induced sudden deaths in the SCDs.

PHT is a condition of increased BPs within the arteries of the lungs. Shortness of breath, fatigue, chest pain, palpitation, swelling of legs and ankles, and cyanosis are common symptoms of PHT. Actually, it is not a diagnosis itself, instead solely a hemodynamic state characterized by resting mean pulmonary artery pressure of 25 mmHg or higher. An increase in pulmonary artery systolic pressure, estimated noninvasively by the echocardiography, helps to identify patients with PHT [48]. The cause is often unknown. The underlying mechanism typically involves inflammation, fibrosis, and subsequent remodelling of the arteries. According to World Health Organization, there are five groups of PHT including pulmonary arterial hypertension, PHT secondary to left heart diseases, PHT secondary to lung diseases, chronic thromboembolic PHT, and PHT with unknown mechanisms [49]. PHT affects about 1% of the world population, and its prevalence may reach 10% above the age of 65 years [50]. Onset is typically seen between 20 and 60 years of age (49). The most common causes are CHD and COPD [49, 51]. The cause of PHT in COPD is generally assumed to be hypoxic pulmonary vasoconstriction leading to permanent medial hypertrophy [52]. But the pulmonary vascular remodeling in the COPD may have a much more complex mechanism than just being the medial hypertrophy secondary to the long-lasting hypoxic vasoconstriction alone [52]. In fact, all layers of the vessel wall appear to be involved with prominent intimal changes [52]. The specific pathological picture could be explained by the combined effects of hypoxia, prolonged stretching of hyperinflated lungs-induced mechanical stress and inflammatory reaction, and the toxic effects of cigarette smoke [52]. On the other hand, PHT is also a common consequence, and its prevalence was detected between 20% and 40% in the SCDs [53, 54]. Whereas we detected the ratio as 12.2% in the present study. The relatively younger mean ages of the study cases (30.8 years of males and 30.3 years of females) may be the cause of the lower prevalence of PHT in the present study. Although the higher prevalences of smoking and alcohol-like atherosclerotic risk factors in male gender, and although the higher prevalences of disseminated teeth losses, ileus, cirrhosis, leg ulcers, digital clubbing, CRD, COPD, and stroke-like atherosclerotic consequences in male gender, and the male gender alone is being a risk factor for the systemic atherosclerosis, the similar prevalences of PHT and ACS in both genders also support nonatherosclerotic backgrounds of them in the SCDs in the present study. Similar to our result, women have up to four times of the risk of men for development of idiopathic PHT, and generally develop symptoms 10 years earlier than men in the literature with the unknown reasons, yet (55). Although COPD and CHD are the most common causes of PHT in the society (51, 56), and although COPD (25.2% vs 7.0%, $p<0.001$) and CHD (18.0% vs 13.2%, $p<0.05$) were higher in male gender in the present study, PHT was not higher in males, again. In another definition, PHT may have a sickled or just hardened RBCs-induced chronic thromboembolic whereas ACS may have an acute thromboembolic backgrounds in the SCDs (57, 58), because the mean age of ACS was lower than PHT (30.3 and 34.0 years, $p<0.05$), but its mortality was much higher than PHT in the literature [43, 44, 49].

COPD is the third leading cause of death with various underlying etiologies all over the world [59, 60]. Aging, physical inactivity, sedentary lifestyle, animal-rich diet, smoking, alcohol, male gender, excess weight, chronic inflammations, prolonged infections, and cancers

may be the major underlying causes. Beside smoking, regular alcohol consumption is also an important risk factor for the pulmonary and systemic atherosclerotic processes, since COPD was one of the most common diagnoses in alcohol dependence [61]. Furthermore, 30-day readmission rates were higher in the COPD patients with alcoholism [62]. Probably an accelerated atherosclerotic process is the main structural background of functional changes seen with the COPD. The inflammatory process of vascular endothelium is enhanced by release of various chemicals by inflammatory cells, and it terminates with an advanced fibrosis, atherosclerosis, and pulmonary losses. COPD may just be the pulmonary consequence of the systemic atherosclerotic process. Since beside the accelerated atherosclerotic process of the pulmonary vasculature, there are several reports about coexistence of associated endothelial inflammation all over the body in COPD [63, 64]. For example, there may be close relationships between COPD, CHD, PAD, and stroke [65], and CHD was the most common cause of deaths in the COPD in a multi-center study of 5.887 smokers [66]. When the hospitalizations were researched, the most common causes were the cardiovascular diseases, again [66]. In another study, 27% of mortality cases were due to the cardiovascular diseases in the moderate and severe COPD [67]. Similarly, COPD may just be the pulmonary consequence of the systemic atherosclerotic process caused by the sickled or just hardened RBCs in the SCDs [59].

Digital clubbing is characterized by the increased normal angle of 165° between nailbed and fold, increased convexity of the nail fold, and thickening of the whole distal finger [68]. Although the exact cause and significance is unknown, the chronic tissue hypoxia is highly suspected [69]. In the previous study, only 40% of clubbing cases turned out to have significant underlying diseases while 60% remained well over the subsequent years [18]. But according to our experiences, digital clubbing is frequently associated with the pulmonary, cardiac, renal, or hepatic diseases or smoking which are characterized by chronic tissue hypoxia (5). As an explanation for that hypothesis, lungs, heart, kidneys, and liver are closely related organs which affect each other's functions in a short period of time. Similarly, digital clubbing is also common in the SCDs, and its prevalence was 10.8% in the present study. It probably shows chronic tissue hypoxia caused by disseminated endothelial damage, inflammation, edema, and fibrosis at the capillaries in the SCDs. Beside the effects of SCDs, smoking, alcohol, cirrhosis, CRD, CHD, and COPD, the higher prevalence of digital clubbing in males (14.8% vs 6.6%, $p < 0.001$) may also show some additional risks of male gender in the systemic atherosclerosis.

Leg ulcers are seen in 10% to 20% of the SCDs, and the ratio was 13.5% in the present study (70). Its prevalence increases with aging, male gender, and SCA (71). Similarly, its ratio was higher in males (19.8% vs 7.0%, $p < 0.001$), and mean age of the leg ulcer patients was higher than the remaining ones in the present study (35.3 vs 29.8 years, $p < 0.000$). The leg ulcers have an intractable nature, and around 97% of them relapse in a period of one year (70). As an evidence of their atherosclerotic background, the leg ulcers occur in the distal segments of the body with a lesser collateral blood supply (70). The sickled or just hardened RBCs-induced chronic endothelial damage, inflammation, edema, and fibrosis at the capillaries may be the major causes, again [71]. Prolonged exposure to the sickled or just hardened bodies due to the pooling of blood in the lower extremities may also explain the leg but not arm ulcers in the SCDs. The sickled or just hardened RBCs-induced venous insufficiencies may also accelerate the highly destructive process by pooling of causative bodies in the legs, and vice versa. Pooling of blood may also have some effects on development of venous ulcers, diabetic ulcers, Buerger's disease, digital clubbing, and onychomycosis in the lower extremities. Furthermore, pooling of blood may be the main cause of delayed wound and fracture healings in the lower extremities. Smoking and alcohol may also have some additional atherosclerotic effects on the leg ulcers in males. Although presence of a continuous damage of hardened RBCs on vascular endothelium, severity of the destructive process is probably

exaggerated by the patients' own immune systems. Similarly, lower WBCs counts were associated with lower crises rates, and if a tissue infarct occurs, lower WBCs counts may decrease severity of pain and tissue damage (31). Because the main action of hydroxyurea may be the suppression of hyperproliferative WBCs and PLTs in the SCDs [72], prolonged resolution of leg ulcers with hydroxyurea may also suggest that the ulcers may be secondary to increased WBCs and PLTs counts-induced exaggerated capillary endothelial inflammation and edema.

Cirrhosis was the 10th leading cause of death for men and the 12th for women in the United States [6]. Although the improvements of health services worldwide, the increased morbidity and mortality of cirrhosis may be explained by prolonged survival of the human being, and increased prevalence of excess weight all over the world. For example, nonalcoholic fatty liver disease (NAFLD) affects up to one third of the world population, and it became the most common cause of chronic liver disease even at childhood, nowadays [73]. NAFLD is a marker of pathological fat deposition combined with a low-grade inflammation which results with hypercoagulability, endothelial dysfunction, and an accelerated atherosclerosis [73]. Beside terminating with cirrhosis, NAFLD is associated with higher overall mortality rates as well as increased prevalences of cardiovascular diseases [74]. Authors reported independent associations between NAFLD and impaired flow-mediated vasodilation and increased mean carotid artery intima-media thickness (CIMT) [75]. NAFLD may be considered as one of the hepatic consequences of the metabolic syndrome and SCDs [76]. Probably smoking also takes role in the inflammatory process of the capillary endothelium in liver, since the systemic inflammatory effects of smoking on endothelial cells is well-known with Buerger's disease and COPD [77]. Increased oxidative stresses, inactivation of antiproteases, and release of proinflammatory mediators may terminate with the systemic atherosclerosis in smokers. The atherosclerotic effects of alcohol is much more prominent in hepatic endothelium probably due to the highest concentrations of its metabolites there. Chronic infectious or inflammatory processes and cancers may also terminate with an accelerated atherosclerosis in whole body [78]. For example, chronic hepatitis C virus (HCV) infection raised CIMT, and normalization of hepatic function with HCV clearance may be secondary to reversal of favourable lipids observed with the chronic infection [78, 79]. As a result, cirrhosis may also be another atherosclerotic consequence of the SCDs.

The increased frequency of CRD can also be explained by aging of the human being, and increased prevalence of excess weight all over the world [80, 81]. Aging, physical inactivity, sedentary lifestyle, animal-rich diet, excess weight, smoking, alcohol, inflammatory or infectious processes, and cancers may be the main underlying causes of the renal endothelial inflammation. The inflammatory process is enhanced by release of various chemicals by lymphocytes to repair the damaged endothelial cells of the renal arteriols. Due to the continuous irritation of the vascular endothelial cells, prominent changes develop in the architecture of the renal tissues with advanced atherosclerosis, tissue hypoxia, and infarcts. Excess weight-induced hyperglycemia, dyslipidemia, elevated BPs, and insulin resistance may cause tissue inflammation and immune cell activation [82]. For example, age ($p = 0.04$), high-sensitivity C-reactive protein ($p = 0.01$), mean arterial BPs ($p = 0.003$), and DM ($p = 0.02$) had significant correlations with the CIMT [81]. Increased renal tubular sodium reabsorption, impaired pressure natriuresis, volume expansion due to the activations of sympathetic nervous system and renin-angiotensin system, and physical compression of kidneys by visceral fat tissue may be some mechanisms of the increased BPs with excess weight [83]. Excess weight also causes renal vasodilation and glomerular hyperfiltration which initially serve as compensatory mechanisms to maintain sodium balance due to the increased tubular reabsorption [83]. However, along with the increased BPs, these changes cause a hemodynamic burden on the kidneys in long term that causes chronic endothelial damage [84]. With prolonged weight excess, there are increased urinary protein excretion, loss of nephron function, and

exacerbated HT. With the development of dyslipidemia and DM in cases with excess weight, CRD progresses much faster [83]. On the other hand, the systemic inflammatory effects of smoking on endothelial cells may also be important in the CRD [85]. Although some authors reported that alcohol was not related with the CRD [85], various metabolites of alcohol circulate even in the renal capillaries, and give harm to the renal capillary endothelium. Chronic inflammatory or infectious processes may also terminate with the accelerated atherosclerosis in the renal vasculature [78]. Although CRD is due to the atherosclerotic process of the renal vasculature, there are close relationships between CRD and other atherosclerotic consequences of the metabolic syndrome including CHD, COPD, PAD, cirrhosis, and stroke [86], and the most common cause of death was the cardiovascular diseases in the CRD again [87]. The sickled or just hardened RBCs-induced capillary endothelial damage may be the main cause of CRD in the SCDs, again [88].

Stroke is an important cause of death, and usually develops as an acute thromboembolic event on the chronic atherosclerotic background. Aging, male gender, smoking, alcohol, and excess weight may be the major underlying causes. Stroke is a common complication of the SCDs, too [89, 90]. Similar to the leg ulcers, stroke is particularly higher with the SCA and higher WBCs counts [91]. Sickling-induced capillary endothelial damage, activations of WBCs, PLTs, and coagulation system, and hemolysis may cause inborn and severe capillary endothelial inflammation, edema, and fibrosis [92]. Probably, stroke may not have a macrovascular origin in the SCDs, and diffuse capillary endothelial inflammation, edema, and fibrosis may be much more important. Infections, inflammations, medical or surgical emergencies, and emotional stresses may precipitate stroke by increasing basal metabolic rate and sickling. A significant reduction of stroke with hydroxyurea may also suggest that a significant proportion of cases is developed secondary to the increased WBCs and PLTs-induced exaggerated capillary endothelial inflammation and edema [45].

The venous capillary endothelium may also be involved in the SCDs [93]. Normally, leg muscles pump veins against the gravity, and the veins have pairs of leaflets of valves to prevent blood from flowing backwards. When the leaflets are damaged, varices and telangiectasias develop. DVT may also cause varicose veins and telangiectasias. Varicose veins are the most common in superficial veins of the legs, which are subject to higher pressure when standing up, thus physical examination must be performed in the upright position. Although the relatively younger mean ages and significantly lower body mass index of the SCDs cases in the literature (10), the prevalences of DVT and/or varices and/or telangiectasias of the lower limbs were relatively higher in the present study (9.0% vs 6.6% in males and females, $p > 0.05$, respectively), indicating an additional venous involvement of the SCDs. Similarly, priapism is the painful erection of penis that can not return to its flaccid state within four hours in the absence of any stimulation [94]. It is an emergency because repeated damaging of the blood vessels may terminate with fibrosis of the corpus cavernosa, a consecutive erectile dysfunction, and eventually a shortened, indurated, and non-erectile penis [94]. It is mainly seen with SCDs, spinal cord lesions (hanging victims), and glucose-6-phosphate dehydrogenase deficiency [95, 96]. Ischemic (veno-occlusive), stuttering (recurrent ischemic), and nonischemic priapisms (arterial) are the three types [97]. Ninety-five percent of clinically presented priapisms are the ischemic (veno-occlusive) disorders in which blood can not return adequately from the penis as in the SCDs, and they are very painful [94, 97]. RBCs support is the treatment of choice in acute whereas hydroxyurea should be the treatment of choice in chronic phases [98]. According to our experiences, hydroxyurea is highly effective for prevention of attacks and consequences of priapism if initiated in early years of life, but it may be difficult due to the excessive fibrosis around the capillaries if initiated later in life.

As a conclusion, the sickled or just hardened RBCs-induced capillary endothelial damage, inflammation, edema, and fibrosis are initiated at birth, and terminate with disseminated tissue hypoxia, multiorgan

failures, and sudden deaths even at childhood. Although RBCs suspensions and corticosteroids in acute and aspirin plus hydroxyurea in acute and chronic phases decrease severity of the inflammatory process, survivals are still shortened in both genders, dramatically. Infections, medical or surgical emergencies, or emotional stresses-induced increased basal metabolic rate aggravates the sickling and capillary endothelial edema, and may terminate with multiorgan failures-induced sudden deaths in the SCDs.

References

1. Widlansky ME, Gokce N, Keane JF Jr, Vita JA. (2003), The clinical implications of endothelial dysfunction. *J Am Coll Cardiol*; 42(7): 1149-1160.
2. Eckel RH, Grundy SM, Zimmet PZ. (2006), **The metabolic syndrome.** *Lancet*; 365(9468): 1415-1428.
3. Franklin SS, Barboza MG, Pio JR, Wong ND. (2006), Blood pressure categories, hypertensive subtypes, and the metabolic syndrome. *J Hypertens*; 24(10): 2009-2016.
4. Third Report of the National Cholesterol Education Program (NCEP) Expert Panel on Detection, Evaluation, and Treatment of High Blood Cholesterol in Adults (Adult Treatment Panel III) final report. *Circulation* 2002; 106(25): 3143-421.
5. Helvacı MR, Aydin LY, Aydin Y. (2012), Digital clubbing may be an indicator of systemic atherosclerosis even at microvascular level. *HealthMED*; 6(12): 3977-3981.
6. Anderson RN, Smith BL. (2003), Deaths: leading causes for 2001. *Natl Vital Stat Rep*; 52(9): 1-85.
7. Helvacı MR, Gokce C, Davran R, Akkucuk S, Ugur M, et al. (2015), Mortal quintet of sickle cell diseases. *Int J Clin Exp Med*; 8(7): 11442-11448.
8. Platt OS, Brambilla DJ, Rosse WF, Milner PF, Castro O, et al. (1994), Mortality in sickle cell disease. Life expectancy and risk factors for early death. *N Engl J Med*; 330(23): 1639-1644.
9. Helvacı MR, Yaprak M, Abyad A, Pocock L. (2018), Atherosclerotic background of hepatosteatosis in sickle cell diseases. *World Family Med*; 16(3): 12-18.
10. Helvacı MR, Kaya H. Effect of sickle cell diseases on height and weight. *Pak J Med Sci* 2011; 27(2): 361-364.
11. Helvacı MR, Aydin Y, Ayyildiz O. (2013). Hydroxyurea may prolong survival of sickle cell patients by decreasing frequency of painful crises. *HealthMED*; 7(8): 2327-2332.
12. Mankad VN, Williams JP, Harpen MD, Mancini E, Longenecker G, et al. (1990). Magnetic resonance imaging of bone marrow in sickle cell disease: clinical, hematologic, and *pathologic correlations.* *Blood*; 75(1): 274-283.
13. Helvacı MR, Aydin Y, Ayyildiz O. (2013). Clinical severity of sickle cell anemia alone and sickle cell diseases with thalassemias. *HealthMED*; 7(7): 2028-2033.
14. Fisher MR, Forfia PR, Chamara E, Houston-Harris T, Champion HC, et al. (2009). Accuracy of Doppler echocardiography in the hemodynamic assessment of pulmonary hypertension. *Am J Respir Crit Care Med*; 179(7): 615-21.
15. Vestbo J, Hurd SS, Agustí AG, Jones PW, Vogelmeier C, Anzueto A, et al. (2013). Global strategy for the diagnosis, management, and prevention of chronic obstructive pulmonary disease: GOLD executive summary. *Am J Respir Crit Care Med*; 187(4): 347-365.
16. Davies SC, Luce PJ, Win AA, Riordan JF, Brozovic M. (1984). Acute chest syndrome in sickle-cell disease. *Lancet*; 1(8367): 36-38.
17. Vandemergel X, Renneboog B. (2008). Prevalence, aetiologies and significance of clubbing in a department of general internal medicine. *Eur J Intern Med*; 19(5): 325-339.
18. Schamroth L. (1976). Personal experience. *S Afr Med J*; 50(9): 297-300.

19. Parfrey NA, Moore W, Hutchins GM. (1985). Is pain crisis a cause of death in sickle cell disease? *Am J Clin Pathol*; 84: 209-212.
20. Helvacı MR, Ayyıldız O, Gundogdu M. (2013). Red blood cell transfusions and survival of sickle cell patients. *HealthMED*; 7(11): 2907-2912.
21. Miller ST, Sleeper LA, Pegelow CH, Enos LE, Wang WC, et al. (2000). Prediction of adverse outcomes in children with sickle cell disease. *N Engl J Med*; 342: 83-89.
22. Helvacı MR, Atci N, Ayyıldız O, Muftuoğlu OE, Pocock L. (2016). Red blood cell supports in severe clinical conditions in sickle cell diseases. *World Family Med*; 14(5): 11-18.
23. Balkaran B, Char G, Morris JS, Thomas PW, Serjeant BE, et al. (1992). Stroke in a cohort of patients with homozygous sickle cell disease. *J Pediatr*; 120: 360-366.
24. Cole TB, Sprinkle RH, Smith SJ, Buchanan GR. (1986). Intravenous narcotic therapy for children with severe sickle cell pain crisis. *Am J Dis Child*; 140: 1255-1259.
25. Yawn BPs, Buchanan GR, Afenyi-Annan AN, Ballas SK, Hassell KL, et al. (2014). Management of sickle cell disease: summary of the 2014 evidence-based report by expert panel members. *JAMA*; 312(10): 1033-1048.
26. Helvacı MR, Ayyıldız O, Gundogdu M. (2014). Hydroxyurea therapy and parameters of health in sickle cell patients. *HealthMED*; 8(4): 451-456.
27. Helvacı MR, Tonyali O, Yaprak M, Abyad A, Pocock L. (2019). Increased sexual performance of sickle cell patients with hydroxyurea. *World Family Med*; 17(4): 28-33.
28. Miller BA, Platt O, Hope S, Dover G, Nathan DG. (1987). Influence of hydroxyurea on fetal hemoglobin production in vitro. *Blood*; 70(6): 1824-1829.
29. Platt OS. (1988). Is there treatment for sickle cell anemia? *N Engl J Med*; 319(22): 1479-1480.
30. Helvacı MR, Aydoğan F, Sevinc A, Camci C, Dilek I. (2014). Platelet and white blood cell count in severity of sickle cell diseases. *Pren Med Argent*; 100(1): 49-56.
31. Charache S. (1997). Mechanism of action of hydroxyurea in the management of sickle cell anemia in adults. *Semin Hematol*; 34(3): 15-21.
32. Charache S, Barton FB, Moore RD, Terrin ML, Steinberg MH, et al. (1996). Hydroxyurea and sickle cell anemia. Clinical utility of a myelosuppressive "switching" agent. The Multicenter Study of Hydroxyurea in Sickle Cell Anemia. *Medicine (Baltimore)*; 75(6): 300-326.
33. Steinberg MH, Barton F, Castro O, Pegelow CH, Ballas SK, et al. (2003). Effect of hydroxyurea on mortality and morbidity in adult sickle cell anemia: risks and benefits up to 9 years of treatment. *JAMA*; 289(13): 1645-1651.
34. Lebensburger JD, Miller ST, Howard TH, Casella JF, Brown RC, et al. (2012). BABY HUG Investigators. Influence of severity of anemia on clinical findings in infants with sickle cell anemia: analyses from the BABY HUG study. *Pediatr Blood Cancer*; 59(4): 675-688.
35. Toghi H, Konno S, Tamura K, Kimura B, Kawano K. (1992). Effects of low-to-high doses of aspirin on platelet aggregability and metabolites of thromboxane A2 and prostacyclin. *Stroke*; 23(10): 1400-1403.
36. Baigent C, Blackwell L, Collins R, Emberson J, Godwin J, et al. (2009). Aspirin in the primary and secondary prevention of vascular disease: collaborative meta-analysis of individual participant data from randomised trials. *Lancet*; 373(9678): 1849-1860.
37. Algra AM, Rothwell PM. (2012). Effects of regular aspirin on long-term cancer incidence and metastasis: a systematic comparison of evidence from observational studies versus randomised trials. *Lancet Oncol*; 13(5): 518-527.
38. Schrör K. (2007). Aspirin and Reye syndrome: a review of the evidence. *Paediatr Drugs*; 9(3): 195-204.
39. Pugliese A, Beltramo T, Torre D. (2008). Reyes and Reye's-like syndromes. *Cell Biochem Funct*; 26(7): 741-746.
40. Hurwitz ES. (1989). Reye's syndrome. *Epidemiol Rev*; 11: 249-253.
41. Macdonald S. (2002). Aspirin use to be banned in under 16-year-olds. *BMJ*; 325(7371): 988.
42. Meremikwu MM, Okomo U. (2011). Sickle cell disease. *BMJ Clin Evid*; 2011: 2402.
43. Poncz M, Kane E, Gill FM. (1985). Acute chest syndrome in sickle cell disease: etiology and clinical correlates. *J Pediatr*; 107(6): 861-866.
44. Sprinkle RH, Cole T, Smith S, Buchanan GR. (1986). Acute chest syndrome in children with sickle cell disease. A retrospective analysis of 100 hospitalized cases. *Am J Pediatr Hematol Oncol*; 8(2): 105-110.
45. Charache S, Terrin ML, Moore RD, Dover GJ, Barton FB, et al. (1995). Effect of hydroxyurea on the frequency of painful crises in sickle cell anemia. Investigators of the Multicenter Study of Hydroxyurea in Sickle Cell Anemia. *N Engl J Med*; 332(20): 1317-1322.
46. Vichinsky E, Williams R, Das M, Earles AN, Lewis N, et al. (1994). Pulmonary fat embolism: a distinct cause of severe acute chest syndrome in sickle cell anemia. *Blood*; 83(11): 3107-3112.
47. Charache S, Scott JC, Charache P. (1979). "Acute chest syndrome" in adults with sickle cell anemia. Microbiology, treatment, and prevention. *Arch Intern Med*; 139(1): 67-69.
48. Gordeuk VR, Castro OL, Machado RF. (2016). Pathophysiology and treatment of pulmonary hypertension in sickle cell disease. *Blood*; 127(7): 820-828.
49. Simonneau G, Gatzoulis MA, Adantia I, Celermajer D, Denton C, et al. (2013). Updated clinical classification of pulmonary hypertension. *J American College Cardiol*; 62(25): 34-41.
50. Hoeper MM, Humbert M, Souza R, Idrees M, Kawut SM, et al. (2016). A global view of pulmonary hypertension. *Lancet Respir Med*; 4(4): 306-322.
51. Naeije R, Barbera JA. (2001). Pulmonary hypertension associated with COPD. *Crit Care*; 5(6): 286-289.
52. Peinado VI, Barbera JA, Abate P, Ramirez J, Roca J, et al. (1999). Inflammatory reaction in pulmonary muscular arteries of patients with mild chronic obstructive pulmonary disease. *Am J Respir Crit Care Med*; 59: 1605-1611.
53. Helvacı MR, Arslanoglu Z, Celikel A, Abyad A, Pocock L. (2018). Pathophysiology of pulmonary hypertension in sickle cell diseases. *Middle East J Intern Med*; 11(2): 14-21.
54. Castro O. (1996). Systemic fat embolism and pulmonary hypertension in sickle cell disease. *Hematol Oncol Clin North Am*; 10(6): 1289-1303.
55. Cunningham CM, Li M, Ruffenach G, Doshi M, Aryan L, et al. (2022). Y-chromosome gene, Uty, protects against pulmonary hypertension by reducing proinflammatory chemokines. *Am J Respir Crit Care Med*; 206(2): 186-196.
56. Duffels MG, Engelfriet PM, Berger RM, van Loon RL, Hoendermis E, Vriend JW, et al. (2007). Pulmonary arterial hypertension in congenital heart disease: an epidemiologic perspective from a Dutch registry. *Int J Cardiol*; 120(2): 198-204.
57. Oudiz RJ. (2016). Classification of pulmonary hypertension. *Cardiol Clin*; 34(3): 359-361.
58. Gladwin MT, Sachdev V, Jison ML, Shizukuda Y, Plehn JF, et al. (2004). Pulmonary hypertension as a risk factor for death in

- patients with sickle cell disease. *N Engl J Med*; 350(9): 886-895.
59. Helvacı MR, Erden ES, Aydin LY. (2013), Atherosclerotic background of chronic obstructive pulmonary disease in sickle cell patients. *HealthMED* ; 7(2): 484-488.
60. Rennard SI, Drummond MB. (2015), Early chronic obstructive pulmonary disease: definition, assessment, and prevention. *Lancet*; 385(9979): 1778-1788.
61. Schoepf D, Heun R. (2015), Alcohol dependence and physical comorbidity: Increased prevalence but reduced relevance of individual comorbidities for hospital-based mortality during a 12.5-year observation period in general hospital admissions in urban North-West England. *Eur Psychiatry* ; 30(4): 459-468.
62. Singh G, Zhang W, Kuo YF, Sharma G. (2016), Association of Psychological Disorders With 30-Day Readmission Rates in Patients With COPD. *Chest*; 149(4): 905-915.
63. Danesh J, Collins R, Appleby P, Peto R. (1998), Association of fibrinogen, C-reactive protein, albumin, or leukocyte count with coronary heart disease: meta-analyses of prospective studies. *JAMA*; 279(18): 1477-1482.
64. Mannino DM, Watt G, Hole D, Gillis C, Hart C, et al. (2006), The natural history of chronic obstructive pulmonary disease. *Eur Respir J*; 27(3): 627-643.
65. Mapel DW, Hurley JS, Frost FJ, Petersen HV, Picchi MA, et al. (2000), Health care utilization in chronic obstructive pulmonary disease. A case-control study in a health maintenance organization. *Arch Intern Med*; 160(17): 2653-2658.
66. Anthonisen NR, Connett JE, Enright PL, Manfreda J. (2002), ; Lung Health Study Research Group. Hospitalizations and mortality in the Lung Health Study. *Am J Respir Crit Care Med* ; 166(3): 333-339.
67. McGarvey LP, John M, Anderson JA, Zvarich M, Wise RA. (2007), TORCH Clinical Endpoint Committee. Ascertainment of cause-specific mortality in COPD: operations of the TORCH Clinical Endpoint Committee. *Thorax*; 62(5): 411-415.
68. Myers KA, Farquhar DR. (2001), The rational clinical examination. Does this patient have clubbing? *JAMA* ; 286(3): 341-347.
69. Toovey OT, Eisenhauer HJ. (2010), A new hypothesis on the mechanism of digital clubbing secondary to pulmonary pathologies. *Med Hypotheses*; 75(6): 511-513.
70. Trent JT, Kirsner RS. (2004), Leg ulcers in sickle cell disease. *Adv Skin Wound Care*: 17(8); 410-416.
71. Minniti CP, Eckman J, Sebastiani P, Steinberg MH, Ballas SK. (2010), Leg ulcers in sickle cell disease. *Am J Hematol*; 85(10): 831-833.
72. Helvacı MR, Aydoğan F, Sevinc A, Camci C, Dilek I. (2014), Platelet and white blood cell counts in severity of sickle cell diseases. *HealthMED* ; 8(4): 477-482.
73. Bhatia LS, Curzen NP, Calder PC, Byrne CD. (2012), Non-alcoholic fatty liver disease: a new and important cardiovascular risk factor? *Eur Heart J*; 33(10): 1190-1200.
74. Pacifico L, Nobili V, Anania C, Verdecchia P, Chiesa C. (2011), Pediatric nonalcoholic fatty liver disease, metabolic syndrome and cardiovascular risk. *World J Gastroenterol*; 17(26): 3082-3091.
75. Mawatari S, Uto H, Tsubouchi H. (2011), Chronic liver disease and arteriosclerosis. *Nihon Rinsho*; 69(1): 153-157.
76. Bugianesi E, Moscatiello S, Ciaravella MF, Marchesini G. (2010), Insulin resistance in nonalcoholic fatty liver disease. *Curr Pharm Des*; 16(17): 1941-1951.
77. Helvacı MR, Aydin LY, Aydin Y. (2012), Chronic obstructive pulmonary disease may be one of the terminal end points of metabolic syndrome. *Pak J Med Sci* ; 28(3): 376-379.
78. Mostafa A, Mohamed MK, Saeed M, Hasan A, Fontanet A, et al. (2010), Hepatitis C infection and clearance: impact on atherosclerosis and cardiometabolic risk factors. *Gut*; 59(8): 1135-1140.
79. Helvacı MR, Ayyıldız O, Gundogdu M, Aydin Y, Abyad A, et al. (2018), Hyperlipoproteinemias may actually be acute phase reactants in the plasma. *World Family Med*; 16(1): 7-10.
80. Levin A, Hemmelgarn B, Culleton B, Tobe S, McFarlane P, et al. (2008), Guidelines for the management of chronic kidney disease. *CMAJ*; 179(11): 1154-1162.
81. Nassiri AA, Hakemi MS, Asadzadeh R, Faizei AM, Alatab S, et al. (2012), Differences in cardiovascular disease risk factors associated with maximum and mean carotid intima-media thickness among hemodialysis patients. *Iran J Kidney Dis*; 6(3): 203-208.
82. Xia M, Guerra N, Sukhova GK, Yang K, Miller CK, et al. (2011), Immune activation resulting from NKG2D/ligand interaction promotes atherosclerosis. *Circulation*; 124(25): 2933-2943.
83. Hall JE, Henegar JR, Dwyer TM, Liu J, da Silva AA, et al. (2004), Is obesity a major cause of chronic kidney disease? *Adv Ren Replace Ther*; 11(1): 41-54.
84. Nerpin E, Ingelsson E, Risérus U, Helmersson-Karlqvist J, Sundström J, Jobs E, et al. (2012), Association between glomerular filtration rate and endothelial function in an elderly community cohort. *Atherosclerosis*; 224(1): 242-246.
85. Stengel B, Tarver-Carr ME, Powe NR, Eberhardt MS, Brancati FL. (2003), Lifestyle factors, obesity and the risk of chronic kidney disease. *Epidemiology*; 14(4): 479-487.
86. Bonora E, Targher G. (2012), Increased risk of cardiovascular disease and chronic kidney disease in NAFLD. *Nat Rev Gastroenterol Hepatol*; 9(7): 372-381.
87. Tonelli M, Wiebe N, Culleton B, House A, Rabbat C, et al. (2006), chronic kidney disease and mortality risk: a systematic review. *J Am Soc Nephrol*; 17(7): 2034-2047.
88. Helvacı MR, Aydin Y, Aydin LY. (2013), Atherosclerotic background of chronic kidney disease in sickle cell patients. *HealthMED*; 7(9): 2532-2537.
89. DeBaun MR, Gordon M, McKinstry RC, Noetzel MJ, White DA, et al. (2014), Controlled trial of transfusions for silent cerebral infarcts in sickle cell anemia. *N Engl J Med*; 371(8): 699-710.
90. Gueguen A, Mahevas M, Nzouakou R, Hosseini H, Habibi A, et al. (2014), Sickle-cell disease stroke throughout life: a retrospective study in an adult referral center. *Am J Hematol*; 89(3): 267-272.
91. Majumdar S, Miller M, Khan M, Gordon C, Forsythe A, et al. (2014), Outcome of overt stroke in sickle cell anaemia, a single institution's experience. *Br J Haematol*; 165(5): 707-713.
92. Kossorotoff M, Grevent D, de Montalembert M. (2014), Cerebral vasculopathy in pediatric sickle-cell anemia. *Arch Pediatr* ; 21(4): 404-414.
93. Helvacı MR, Gokce C, Sahan M, Hakimoglu S, Coskun M, et al. (2016), Venous involvement in sickle cell diseases. *Int J Clin Exp Med* ; 9(6): 11950-11957.
94. Kaminsky A, Sperling H. (2015), Diagnosis and management of priapism. *Urologe A* ; 54(5): 654-661.
95. Anele UA, Le BV, Resar LM, Burnett AL. (2015), How I treat priapism. *Blood* ; 125(23): 3551-3558.
96. Bartolucci P, Lionnet F. (2014), Chronic complications of sickle cell disease. *Rev Prat*; 64(8): 1120-1126.
97. Broderick GA. (2012), Priapism and sickle-cell anemia: diagnosis and nonsurgical therapy. *J Sex Med* ; 9(1): 88-103.

98. Ballas SK, Lyon D. (2016), Safety and efficacy of blood exchange transfusion for priapism complicating sickle cell disease. *J Clin Apher*; 31(1): 5-10.



This work is licensed under Creative Commons Attribution 4.0 License

To Submit Your Article Click Here: **Submit Manuscript**

DOI: [10.31579/2641-5194/065](https://doi.org/10.31579/2641-5194/065)

Ready to submit your research? Choose Auctores and benefit from:

- fast, convenient online submission
- rigorous peer review by experienced research in your field
- rapid publication on acceptance
- authors retain copyrights
- unique DOI for all articles
- immediate, unrestricted online access

At Auctores, research is always in progress.

Learn more <https://auctoresonline.org/journals/gastroenterology-pancreatology-and-hepatobiliary-disorders>