

# Viruses In Beef, Mutton, Chevron, Venison, Fish and Poultry Meat Products

Fahim A. Shaltout

Food Control Department, Faculty of Veterinary Medicine, Benha university, Egypt.

**\*Corresponding Author:** Fahim A. Shaltout, Food Control Department, Faculty of Veterinary Medicine, Benha university, Egypt.

**Received date:** December 12, 2023; **Accepted date:** December 27, 2023; **Published date:** January 03, 2024

**Citation:** Fahim A. Shaltout. (2024), Viruses in Beef, Mutton, Chevron, Venison, Fish and Poultry Meat Products., *J. Women Health Care and Issues*, 7(1): DOI:10.31579/2637-8892/177

**Copyright:** © 2024, Fahim A. Shaltout. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

## Abstract

Beef means meat of cattle, mutton means meat of sheep, chevon means meat of flesh of the goat used as food, venison means meat of deer, fish means meat of fish and shellfish, poultry means chicken, duck, geese, turkey, pigeon and rabbit. Beef, mutton, chevon, venison, fish and poultry meat act as a main sources of animal protein, in recent decades, viruses have been increasingly known as important causes of foodborne diseases mostly due to the improved methods of diagnosis and investigation of viruses. Viruses' transmission through consumption of infected beef, mutton, chevon, venison, fish and poultry meat products or contact with contaminated beef, mutton, chevon, venison, fish and poultry meat products and water is now well known. The viruses most frequently involved in foodborne infections are public noroviruses, hepatitis A virus, human rotavirus, and hepatitis E virus. Beef, mutton, chevon, venison, fish and poultry meat act as major sources of animal protein, NoV and RV infections are common cause of acute human gastroenteritis, while hepatitis A virus and hepatitis E virus cause human hepatitis worldwide. Most of the cases remain unreported due to subclinical cases.

**Key words:** virus; beef; mutton; chevon; venison; fish; poultry meat products

## Introduction

Beef, mutton, chevon, venison, fish and poultry meat act as a main sources of animal protein, Foodborne viruses are generally very infectious and their spreading are rapidly from one individual to the next, although several exceptions are existed as hepatitis E virus. The most of foodborne virus's outbreaks were linked with the infected food handlers, as hepatitis A virus are mainly transmitted between humans [133,136,137, 1,2,3 and 4]. In contrast hepatitis E virus has been identified as an important disease. Beef, mutton, chevon, venison, fish and poultry meat can potentially be contaminated throughout the whole food product chain and sources of contaminations can include equipment, other contaminated food and Beef, mutton, chevon, venison, fish and poultry meat or meat products, originating from infected animals and water [131,135,139, 5,6,7,8 and 9]. Beef, mutton, chevon, venison, fish and poultry meat act as a main sources of animal protein, and shellfish are the major food categories involved in foodborne of viral gastroenteritis origin [140,141,142, 10,11,12,13 and 14]. However, risky Beef, mutton, chevon, venison, fish and poultry meat are considered particularly those that are intended for direct consumption or that are not properly heat treated before consumption [135,136,137,15,16,17,18 1nd 19]. investigation of viruses in Beef, mutton, chevon, venison, fish and poultry meat products [131,132,133,134,135 and 136]

Polymerase Chain Reaction method for Hepatitis A virus investigation

PCR method, polymerase chain reaction method, is a mean for amplification of a region of DNA whose arrangement is known or lies between two portions of known arrangement.

Before PCR, DNA of interest could be amplified by over-expression in cells and this with limited yield. Components

- DNA template
- Primers
- Enzyme
- dNTPs
- Mg<sup>2+</sup>
- buffers1- DNA template
- 1- DNA template
- DNA containing region to be arrangement
- Size of target DNA to be amplified: up to 3 Kb
- 2- Primers
- Two sets of primers

- Generally 20-30 nucleotides long
  - Synthetically produced complimentary to the 3' ends of target DNA
- not complimentary to each other Primers (• Not containing inverted repeat arrangement to avoid formation of internal structures
- 40-60% GC content preferred for better annealing
  - $T_m$  of primers can be calculated to determine annealing  $T_0$
  - $T_m = .41(\%G+C) + 16.6\log(J+) + 81.5$  where  $J+$  is the concentration of monovalent ions

### 3-Enzyme

- Usually Taq Polymerase or anyone of the natural or Recombinant thermostable polymerases.
- Stable at  $T_0$  up to 950 C
- High processivity
- Taq Pol has 5'-3' exo only, no proofreading

The PCR Cycle Comprised of 3 steps: -

- 1- Denaturation of DNA at 95 C
- 2- Primer hybridization (annealing) at 40-50 C
- 3- DNA synthesis (Primer extension) at 72 c72

### RT-PCR method

- Reverse Transcriptase PCR
- Uses RNA as the initial template
- RNA-directed DNA polymerase (rTh)
- Yields ds cDNA

Investigation of amplification products

- Gel electrophoresis
- Sequencing of amplified fragment
- Southern blot

### Advantages

- Automated, fast, reliable (reproducible) results
- Contained :(less chances of contamination)
- High output
- Sensitive
- Broad uses
- Defined, easy to follow protocols

ELISA method for Food Borne Viruses investigation:

ELISA method Is a biochemical method used mainly in immunology to detect the presence or absence of an antibody or an antigen in a beef, mutton, chevon, venison, fish and poultry meat samples.

The method is divided into

- 1- Competitive ELISA method.
- 2- Sandwich ELISA method or direct ELISA method.
- 3- Indirect ELISA method.

### Competitive ELISA method

- The labeled antigen competes for primary antibody binding places with the beef, mutton, chevon, venison, fish and poultry meat samples antigen. The more antigen in the beef, mutton, chevon, venison, fish and poultry meat samples, the less labelled antigen is retained in the well and the weaker the signal.

### Sandwich ELISA

- The ELISA plate is coated with Antibody to detect specific antigen

Prepare a surface to which a known quantity of capture antibody is bound.

- Block any nonspecific binding sites on the surface
- Apply the antigen-containing sample to the plate.

Wash the plate, so the unbound antigen is removed.

Apply enzyme linked primary antibodies as investigation antibodies which also bind specifically to the antigen.

Wash the plate, so the the unbound antibody enzyme conjugates are removed.

- Apply a chemical which is converted by the enzyme into a coloured product.

- Detect the absorbency of the plate wells to investigate the presence and quantity of antigen

### Indirect ELISA

- The protein antigen to be tested is added to each well of ELISA plate, where it is given time to adhere to the plastic by charge interactions.

- A solution of non-reacting protein is added to block any plastic surface in the well that remains uncoated by the protein antigen

Then the serum is added, which contains a mixture of the serum antibodies, of unknown concentration, some of which may bind specifically to the test antigen that is coating the well. Afterwards, a secondary antibody is added, which will bind to the antibody bound to the test antigen in the well. This secondary antibody often has an enzyme attached to it

A substrate for this enzyme is then added. This substrate changes colour upon reaction with the enzyme. The colour change shows that secondary antibody has bound to primary antibody, which strongly implies that the donor has had an immune reaction to the test antigen. The higher the concentration of the primary antibody that was presents in the serum, the stronger the colour change. Often a spectrometer is used to give quantitative values for colour strength

### Discussion:

Beef, mutton, chevon, venison, fish and poultry meat act as a main sources of animal protein. Viruses are transmitted through foods in the form of extremely small particles, they ranging in size nearly from twenty-five nanometers to less than one hundred nanometers in diameter [133,134,135,20,21,222,23,24 and 25]. Virus structure is mainly nucleic acid core with a protein coat. A few have an additional, lipid-containing envelope (140,141,142,26,27,28,29 and 30). The particles are roughly spherical in shape and are totally inert, in the sense that they cannot carry out any of what are commonly regarded as life processes [131,132,133,31,32,33,34 and 35].

Beef, mutton, chevon, venison, fish and poultry meat act as a main sources of animal protein, viruses are of concern to health because of their ability to produce infections, some of which result in disease [141,142,140, 36,38,38,39 and 40]. They do this depend up on a very selective basis. Viruses that infect public tend not to be capable of infecting other species, with the exception of our closest evolutionary relatives [144,143,139,41,42,43,44 and 45]. Viruses that infect other animal species

tend not to be infectious for humans [46,47,48 and 49]. The exceptions, viruses that are occasionally transmitted from Beef, mutton, chevon, venison, fish and poultry meat act as a main sources of animal protein, to man, are not known to be transmitted through Beef, mutton, chevon, venison, fish and poultry meat and meat products [131,138,139,50,51,52,53,54 and 55]. In addition to their species specificity, viruses show a distinct individual preference for infecting certain tissues or organs of the host's body [144,143,142,141,56,57,58,59 and 60]. This tissue specificity determines which cells of the host's body become infected and what symptoms are likely to disease result from virus infection. Whether or not they cause disease, virus infections tend to be self-limiting [144,140,137,61,62,63,64 and 65].

Beef, mutton, chevon, venison, fish and poultry meat act as a main sources of animal protein, The body's immune processes ordinarily suppress a virus infection after some period of time, so the presence of viruses as obligate parasites depend up on their ability to pass from one host to another host. Viruses that infect humans are principally transmitted directly from person to person, either by actual touching or by aerosols over short distances. However, they are also capable of being transmitted indirectly through food and water, as well as a few other means [144,140,139,66,67,68,69 and 70].

Beef, mutton, chevon, venison, fish and poultry meat act as a main sources of animal protein, Virus contamination of foods has been categorized as primary or secondary, depending upon whether the viruses are present in the beef, mutton, chevon, venison, fish and poultry at the time of slaughter. In the case of Beef, mutton, chevon, venison, fish and poultry meat and meat products (137,138,139,71,72,73,74,75 and 76), the viruses that are already present at the time of slaughter are of little concern to public health. Instead, the outbreaks recorded indicated that what problems were lied in beef, mutton, chevon, venison, fish and poultry meat contamination, usually mishandling of Beef, mutton, chevon, venison, fish and poultry meat by a person with an gut virus infection (144,143,77,78,79,80 and 81). public gut viruses in sewage have also contaminated Beef, mutton, chevon, venison, fish and poultry meat and meat products; but neither insects nor rodents are known to have served as vectors in secondary contamination of Beef, mutton, chevon, venison, fish and poultry meat, despite the obvious possibility that they might do so [135,136,82,83,84,85 and 86].

Contamination of Beef, mutton, chevon, venison, fish and poultry meat act as a main sources of animal protein does not guarantee that a consumer infection will result (nor, for that matter, do most virus infections result in overt disease). Virus that has been introduced into Beef, mutton, chevon, venison, fish and poultry meat cannot possibly multiply, but may be inactivated (deprived of its infectivity) before the Beef, mutton, chevon, venison, fish and poultry meat are eaten[87,88,89,90 and 91]. This can come about in a number of ways, the one of most practical significance being thermal processing or cooking (92,93,94,95,96 and 97). The times and temperatures required for virus inactivation in Beef, mutton, chevon, venison, fish and poultry meat cannot be specified precisely [131,132,133,98,99,100,101,102 and 103].

Beef, mutton, chevon, venison, fish and poultry meat act as a main sources of animal protein, Viruses in a rare steak probably are no threat to public health because viruses within the muscle are likely to be of animal origin and therefore not infectious for the consumer[104,105,106,107 and 108]. Viruses in ground beef, however, may be of human origin: the heat stability of viral contaminant varies with the fat content of the ground beef, but complete inactivation can apparently be assured by cooking the Beef, mutton, chevon, venison, fish and poultry meat until all pink colour disappears from the center[144,143,109,110,111,112,113 and 114]. Virus on the surface of Beef, mutton, chevon, venison, fish and poultry meat can probably be inactivated by ultraviolet light, and ionizing radiation can inactivate virus in subsurface locations [115,116,117,118 and 119]. Although the coat proteins of some viruses are apparently biodegradable, microbial decomposition of Beef, mutton, chevon, venison, fish and poultry meat through prolonged storage evidently has little effect upon the virus [133, 134,120,121,122,123 and 124].

Beef, mutton, chevon, venison, fish and poultry meat act as a main sources of animal protein, There is one important exception to some of the above generalizations that should be mentioned[131,132,125,126,127,128,129 and 130] the virus of foot and mouth disease, which is no direct threat to human health but has great economic significance, is chemically degraded in voluntary muscle by the acid of rigor mortis but is protected from this, and withstands a great deal of heat, in lymph nodes, bone marrow, and large blood clots[143,142,140 and 144].

Beef, mutton, chevon, venison, fish and poultry meat act as a main sources of animal protein, Many kinds of viruses in Beef, mutton, chevon, venison, fish and poultry meat can be detected on the basis of their ability to produce infections in cell cultures. The absolutely necessary steps in the investigation process are to make a fluid suspension of the sample and inoculate it into a culture of susceptible cells; however, in practice, several additional steps are usually required. Detection of viruses that are of significance to public health, but no type of cell culture is known to be susceptible to the virus of hepatitis A or to some of the viral gastroenteritis agents. The methods that are available are used, despite their cost and complexity, because they are not valid indicators, the presence of which would indicate the virus contamination of beef, mutton, chevon, venison, fish and poultry meat had occurred [143,142,140 and 139]. Plant or market samples of Beef, mutton, chevon, venison, fish and poultry meat and meat products have been tested for viral contaminants. Ground beef has attracted a great attention, human viruses were detected in market Beef, mutton, chevon, venison, fish and poultry meat and meat products, Hepatitis has shown that viral contamination of ground beef can be a threat to public health. Gut virus infections are common in slaughter animals. Viruses were also found in some by-products, but the viruses apparently were not infectious for human, and none were found in market Beef, mutton, chevon, venison, fish and poultry meat and meat products [135,136,137 and 138]. human viral diseases associated with Beef, mutton, chevon, venison, fish and poultry meat have included only hepatitis A, a lingering, debilitating disease that is very specific for human and is transmitted by a fecal-oral cycle. Other human gut viruses might well be transmitted through Beef, mutton, chevon, venison, fish and poultry meat in the same way on occasion, as is beginning to be observed with other foods. They are not transmitted between humans and animals and that, where this could be determined, all of the events of contamination that led to outbreaks took place in Beef, mutton, chevon, venison, fish and poultry meat and meat products service or retail establishments [131,132,133 and 134]. Infected butcher contaminated steak tartare (seasoned raw ground beef) in such a way that consumers became ill with hepatitis A., contaminated Beef, mutton, chevon, venison, fish and poultry meat and meat products cause more consumer illnesses, a cafeteria , contaminated roast meat during boning and slicing sufficiently to cause illness in students and faculty. In each of these instances, the virus that contaminated the Beef, mutton, chevon, venison, fish and poultry meat and meat products originated in the human gut: contamination was either direct, or indirect by way of wastewater [33, 138,137,136 and 135].

### Conclusion:

Beef, mutton, chevon, venison, fish and poultry meat -associated viral disease reveals that are not transmitted to consumers causative beef, mutton, chevon, venison, fish and poultry meat. However, viruses that originate in the human gut are as likely to contaminate Beef, mutton, chevon, venison, fish and poultry meat as other foods and, if not inactivated before the Beef, mutton, chevon, venison, fish and poultry meat is eaten, may cause infections in consumers. Recorded incidents have resulted from mishandling Beef, mutton, chevon, venison, fish and poultry meat in food service or retailing, rather than in slaughtering or processing. Viral contamination of Beef, mutton, chevon, venison, fish and poultry meat can be avoided by the same precautions in sanitary Beef, mutton, chevon, venison, fish and poultry meat and meat products handling that are applicable to any other foods. From the standpoint of public health, the viral hazards associated with Beef, mutton,

chevon, venison, fish and poultry meat is significant, but by no means as severe as those of botulism or salmonellosis.

## References:

1. Abd Elaziz, O., Fatin S. Hassanin, Fahim A. Shaltout and Othman A. Mohamed (2021 a): Prevalence of Some Foodborne Parasitic Affection in Slaughtered Animals in Local Egyptian Abattoir. *Journal of Nutrition Food Science and Technology* 2(3): 1-5.
2. Abd Elaziz, O., Fatin, S Hassanin , Fahim, A Shaltout, Othman, A Mohamed (2021 b): Prevalence of some zoonotic parasitic affections in sheep carcasses in a local abattoir in Cairo, Egypt. *Advances in Nutrition & Food Science* 6(2): 6(2): 25-31.
3. Al Shorman,A.A.M. ;Shaltout,F.A. and hilat,N (1999):Detection of certain hormone residues in meat marketed in Jordan. *Jordan University of Science and Technology, 1st International Conference on Sheep and goat Diseases and Productivity*, 23-25 .
4. Harris L.J., Farber J.N., Beuchat L.R., Parish M.E., Suslow T.V., Garrett E.H., Busta F.F. (2006):Outbreaks associated with fresh produce: incidence, growth, and survival of pathogens in fresh and fresh-cut produce. *Compr. Rev. Food Sci. Food Saf.* ; 2:78–141.
5. Ebeed Saleh , Fahim Shaltout , Essam Abd Elaal (2021); Effect of some organic acids on microbial quality of dressed cattle carcasses in Damietta abattoirs, Egypt. *Damanhour Journal of Veterinary Sciences* 5(2): 17-20.
6. Edris A, Hassanin, F. S; Shaltout, F.A., Azza H Elbaba and Nairoz M Adel(2017): Microbiological Evaluation of Some Heat Treated Fish Products in Egyptian Markets. *EC Nutrition* 12.3 (2017): 124-132.
7. Edris ,A., Hassan,M.A., Shaltout,F.A. and Elhosseiny , S(2013): Chemical evaluation of cattle and camel meat. *BENHA VETERINARY MEDICAL JOURNAL*, 24( 2): 191-197
8. Edris ,A.M., Hassan,M.A., Shaltout,F.A. and Elhosseiny , S(2012): Detection of E.coli and Salmonella organisms in cattle and camel meat. *BENHA VETERINARY MEDICAL JOURNAL*, 24( 2): 198-204
9. Edris A.M.; Hemmat M. I., Shaltout F.A.; Elshater M.A., Eman F.M.I. (2012): STUDY ON INCIPIENT SPOILAGE OF CHILLED CHICKEN CUTS-UP. *BENHA VETERINARY MEDICAL JOURNAL*, VOL. 23, NO. 1, JUNE 2012: 81-86
10. Edris A.M.; Hemmat M.I.; Shaltout F.A.; Elshater M.A., Eman, F.M.I.(2012):CHEMICAL ANALYSIS OF CHICKEN MEAT WITH RELATION TO ITS QUALITY. *BENHA VETERINARY MEDICAL JOURNAL*, 23( 1): 87-92
11. Edris, A.M.; Shaltout, F.A. and Abd Allah, A.M. (2005): Incidence of Bacillus cereus in some meat products and the effect of cooking on its survival . *Zag. Vet. J.*33 (2):118-124
12. Edris, A.M.; Shaltout, F.A. and Arab, W.S. (2005): Bacterial Evaluation of Quail Meat.*Benha Vet. Med.J.*16 (1):1-14.
13. Le Guyader F.S., Bon F., DeMedici D., Parnaudeau S., Bertone A., Crudeli S., Doyle A., Zidane M., Suffredini E., Kohli E., Maddalo F., Monini M., Gallay A., Pommepuy M., Pothier P., Ruggeri F.M. (2006):Detection of multiple noroviruses associated with an international gastroenteritis outbreak linked to oyster consumption. *J. Clin. Microbiol.* ;44:3878–3882.
14. Edris, M. A.; Hassanin, F. S; Shaltout, F.A; Azza, H.ELbaba; Nairoz, M. Adel. (2017): Microbiological evaluation of some frozen and salted fish products in Egyptian markets. *BENHA VETERINARY MEDICAL JOURNAL*, 33, (2): 317-328
15. Edris, A.M.; Shaltout, F.A. ;Salem, G.H. and El-Toukhy,E.I. (2011): Incidence and isolation of Salmonellae from some meat products. *Benha University ,Faculty of Veterinary Medicine , Fourth Scientific Conference 25-27th May 2011Veterinary Medicine and Food Safety )* 172-179 benha , Egypt..
16. Leroy E.M., Epelboin A., Mondonge V., Pourrut H., Gonzalez J.P., Muyembe-Tamfum J.J., Formenty P. (2009): public Ebola outbreak resulting from direct exposure to fruit bats in Luebo, Democratic Republic of Congo, 2007. *Vector Borne Zoonotic Dis.* ;9(6):723–728.
17. Edris, A.M.; Shaltout, F.A. ;Salem, G.H. and El-Toukhy,E.I. (2011): Plasmid profile analysis of Salmonellae isolated from some meat products.*Benha University ,Faculty of Veterinary Medicine , Fourth Scientific Conference 25-27th May 2011Veterinary Medicine and Food Safety )*194-201 benha , Egypt..
18. Ragab A , Abobakr M. Edris, Fahim A.E. Shaltout, Amani M. Salem(2022): Effect of titanium dioxide nanoparticles and thyme essential oil on the quality of the chicken fillet. *BENHA VETERINARY MEDICAL JOURNAL*41(2): 38-40.
19. Hassan, M.A, Shaltout, F. A, Arfa M.M , Mansour A.H and Saudi, K. R(2013): BIOCHEMICAL STUDIES ON RABBIT MEAT RELATED TO SOME DISEASES. *BENHA VETERINARY MEDICAL JOURNAL* 25(1):88-93.
20. Hassan, M.A and Shaltout, F.A. (1997): Occurrence of Some Food Poisoning Microorganisms In Rabbit Carcasses . *Alex.J.Vet.Science*, 13(1):55-61.
21. Hassan M, Shaltout FA\* and Saqur N (2020): Histamine in Some Fish Products. *Archives of Animal Husbandry & Dairy Science* 2(1): 1-3.
22. Hassan, M. A , Fahim A. Shaltout , Nabila E. El-sheikh , Naglaa M. Sakr(2019): Assessment of histamine residues in smoked and salted fish. *Benha Veterinary Medical Journal* 37,2 (2019) 50-52
23. Hassan, M.A and Shaltout, F.A. (2004): Comparative Study on Storage Stability of Beef, Chicken meat, and Fish at Chilling Temperature. *Alex.J.Vet.Science*, 20(21):21-30.
24. Hassan, M.A ; Shaltout, F.A. ; Arafa ,M.M. ; Mansour , A.H. and Saudi , K.R.(2013): Biochemical studies on rabbit meat related to sime diseases . *Benha Vet. Med.J.*25 (1):88-93
25. Hassan, M.A ; Shaltout, F.A. ; Maarouf , A.A. and El-Shafey, W.S.(2014): Psychrotrophic bacteria in frozen fish with special reference to pseudomonas species .*Benha Vet. Med.J.*27 (1):78-83
26. Hassan, M.A ; Shaltout, F.A. ; Arafa ,M.M. ; Mansour , A.H. and Saudi , K.R.(2013): Bacteriological studies on rabbit meat related to sime diseases *Benha Vet. Med.J.*25 (1):94-99.
27. Hassanin, F. S; Hassan,M.A., Shaltout, F.A., Nahla A. Shawqy and 2Ghada A. Abd-Elhameed(2017): Chemical criteria of chicken meat. *BENHA VETERINARY MEDICAL JOURNAL*, 33(2):457-464.
28. Hassanin, F. S; Hassan,M.A.; Shaltout, F.A. and Elrais-Amina, M(2014): CLOSTRIDIUM PERFRINGENS IN VACUUM PACKAGED MEAT PRODUCTS. *BENHA VETERINARY MEDICAL JOURNAL*, 26(1):49-53.
29. Hassanien, F.S. ; Shaltout, F.A.; Fahmey, M.Z. and Elsukkary, H.F.(2020): Bacteriological quality guides in local and imported beef and their relation to public health. *Benha Veterinary Medical Journal* 39: 125-129
30. Li W., She R., Wei H., Zhao J., Wang Y., Sun Q., Zhang Y., Wang D., Li R. (2009):Prevalence of hepatitis E virus in swine under different breeding environment and abattoir in Beijing, China. *Vet. Microbiol.* ;133:75–83.

31. 31.Hassanin, F. S; Shaltout,F.A. and , Mostafa E.M(2013): Parasitic affections in edible offal. *Benha Vet. Med.J.25 (2):*34-39.
32. 32.Hassanin, F. S; Shaltout, F.A., Lamada, H.M., Abd Allah, E.M.(2011): THE EFFECT OF PRESERVATIVE (NISIN) ON THE SURVIVAL OF LISTERIA MONOCYTOGENES. *BENHA VETERINARY MEDICAL JOURNAL (2011)-SPECIAL ISSUE [I]:* 141-145
33. 33.Khattab, E.,Fahim Shaltout and Islam Sabik (2021): Hepatitis A virus related to foods. *BENHA VETERINARY MEDICAL JOURNAL 40(1):* 174-179.
34. 34.Saad M. Saad , Fahim A. Shaltout , Amal A. A. Farag & Hashim F. Mohammed (2022): Organophosphorus Residues in Fish in Rural Areas. *Journal of Progress in Engineering and Physical Science 1(1):* 27-31.
35. 35.Saif,M. , Saad S.M. , Hassanin, F. S; Shaltout FA, Marionette Zaghloul (2019): Molecular detection of enterotoxigenic *Staphylococcus aureus* in ready-to-eat beef products. *Benha Veterinary Medical Journal 37 (2019) 7-11*
36. 36.Saif,M. , Saad S.M. , Hassanin, F. S; Shaltout, F.A., Marionette Zaghloul(2019): Prevalence of methicillin-resistant *Staphylococcus aureus* in some ready-to-eat meat products. *Benha Veterinary Medical Journal 37 (2019) 12-15*
37. 37.Farag, A. A., Saad M. Saad<sup>1</sup>, Fahim A. Shaltout<sup>1</sup>, Hashim F. Mohammed(2023 a): Studies on Pesticides Residues in Fish in Menofia Governorate. *Benha Journal of Applied Sciences , 8(5):* 323-330.
38. 38.Farag, A. A., Saad M. Saad<sup>1</sup>, Fahim A. Shaltout<sup>1</sup>, Hashim F. Mohammed(2023 b): Organochlorine Residues in Fish in Rural Areas. *Benha Journal of Applied Sciences , 8 (5):* 331-336.
39. 39.Shaltout, F.A., Mona N. Hussein, Nada Kh. Elsayed (2023): Histological Detection of Unauthorized Herbal and Animal Contents in Some Meat Products. *Journal of Advanced Veterinary Research 13(2):* 157-160.
40. 40.Shaltout, F. A. , Heikal, G. I. , Ghanem, A. M.(2022): Mycological quality of some chicken meat cuts in Gharbiya governorate with special reference to *Aspergillus flavus* virulent factors. *benha veteriv medical journal veterinary 42(1):* 12-16
41. 41.Shaltout, F.A., Ramadan M. Salem, Eman M. Eldiasty, Fatma A. Diab (2022): Seasonal Impact on the Prevalence of Yeast Contamination of Chicken Meat Products and Edible Giblets. *Journal of Advanced Veterinary Research 12(5):* 641-644
42. 42.Shaltout, F.A., Abdelazez Ahmed Helmy Barr and Mohamed Elsayed Abdelaziz (2022): Pathogenic Microorganisms in Meat Products. *Biomedical Journal of Scientific & Technical Research 41(4):* 32836-32843.
43. 43.Shaltout, F.A., Thabet, M.G. and Koura, H.A. (2017). Impact of Some Essential Oils on the Quality Aspect and Shelf Life of Meat. *J Nutr Food Sci., 7: 647*
44. 44.Shaltout, F.A.,, Islam Z. Mohammed<sup>2</sup>, El -Sayed A. Afify(2020): Bacteriological profile of some raw chicken meat cuts in Ismailia city, Egypt. *Benha Veterinary Medical Journal 39 (2020) 11-15*
45. 45.Shaltout, F.A.,Islam, Z. Mohammed<sup>2</sup>, El -Sayed A. Afify(2020): Detection of *E. coli O157* and *Salmonella* species in some raw chicken meat cuts in Ismailia province, Egypt. *Benha Veterinary Medical Journal 39 (2020) 101-104*
46. 46.Shaltout, F.A., E.M. El-diasty and M. A. Asmaa- Hassan (2020): HYGIENIC QUALITY OF READY TO EAT COOKED MEAT IN RESTAURANTS AT Cairo. *Journal of Global Biosciences 8(12):* 6627-6641.
47. 47.Shaltout, F.A., Marrionet Z. Nasief , L. M. Lotfy , Bossi T. Gamil(2019 a): Microbiological status of chicken cuts and its products. *Benha Veterinary Medical Journal 37 (2019) 57-63*
48. 48.Shaltout, F.A.(2019): Poultry Meat. *Scholarly Journal of Food and Nutrition 22 1-2.*
49. 49.Shaltout, F.A.(2019): Food Hygiene and Control. *Food Science and Nutrition Technology 4(5):* 1-2
50. 50.Hassanin, F. S; Shaltout, F.A., Seham N. Homouda and Safaa M. Araakeb(2019 b): Natural preservatives in raw chicken meat. *Benha Veterinary Medical Journal 37 (2019) 41-45*
51. 51.Lopman B.A., van Duynhoven Y., Hanon F.X., Reacher M., Koopmans M., Brown D. (2002):Consortium on food-borne viruses in Europe: laboratory capability in Europe for food-borne viruses. *Euro Surveill. ;7(4):*61-65.
52. 52.Hazaa,W. , Shaltout, F.A., Mohamed El-Shater (2019 c): Prevalence of some chemical hazards in some meat products. *Benha Veterinary Medical Journal 37 (2) 32-36*
53. 53.Hazaa,W, Shaltout, F.A., Mohamed El-Shater(2019 d): Identification of Some Biological Hazards in Some Meat Products. *Benha Veterinary Medical Journal 37 (2) 27-31*
54. 54.Gaafar,R. , Hassanin, F. S; Shaltout, F.A., Marionette Zaghloul (2019 a): Molecular detection of enterotoxigenic *Staphylococcus aureus* in some ready to eat meat-based sandwiches. *Benha Veterinary Medical Journal 37 (2) 22-26*
55. 55.Gaafar,R. , Hassanin, F. S; Shaltout, F.A., Marionette Zaghloul( 2019 b): Hygienic profile of some ready to eat meat product sandwiches sold in Benha city, Qalubiyah Governorate, Egypt. *Benha Veterinary Medical Journal 37 (2) 16-21*
56. 56.Saad S.M. , Shaltout, F.A., Nahla A Abou Elroos, Saber B El-nahas( 2019 a) : Antimicrobial Effect of Some Essential Oils on Some Pathogenic Bacteria in Minced Meat. *J Food Sci Nutr Res. 2019; 2 (1):* 012-020.
57. 57.Saad S.M. , Shaltout, F.A., Nahla A Abou Elroos<sup>2</sup> and Saber B El-nahas( 2019 b): Incidence of *Staphylococci* and *E. coli* in Meat and Some Meat Products. *EC Nutrition 14.6 (2019).*
58. 58.Saad S.M. , Hassanin, F. S ; Shaltout, F.A., Marionette Z Nassif, Marwa Z Seif.(2019 c): Prevalence of Methicillin-Resistant *Staphylococcus Aureus* in Some Ready-to-Eat Meat Products. *American Journal of Biomedical Science & Research 4(6):*460-464.
59. 59.Shaltout, Fahim(2019): Pollution of Chicken Meat and Its Products by Heavy Metals.
60. 60.Research and Reviews on Healthcare: Open Access Journal, 4, 3(381-3382).
61. 61.Shaltout, F. A.; E.M EL-diasty; M. S. M Mohamed (2018): Effects of chitosan on quality attributes fresh meat slices stored at 4 C. *BENHA VETERINARY MEDICAL JOURNAL, VOL. 35, NO. 2: 157-168.*
62. 62.Shaltout and Abdel-Aziz, 2004 *Salmonella enterica* serovar Enteritidis in poultry meat and their epidemiology *Vet. Med. J. Giza, 52 (2004), pp. 429-436*
63. 63.Shaltout, F.A., Hala F El-Shorah, Dina I El Zahaby, Lamiaa M Lotfy(2018):Bacteriological Profile of Chicken Meat Products. *SciFed Food & Dairy Technology Journal, 2:*
64. 64.Shaltout, F.A., Mohamed, A.H. El-Shater ., Wafaa Mohamed Abd El-Aziz(2015): Bacteriological assessment of Street Vended Meat Products sandwiches in kalyobia Governorate. *BENHA VETERINARY MEDICAL JOURNAL, 28( 2):*58-66,
65. 65.Shaltout, F.A., Mohamed A El shatter and Heba M Fahim(2019): Studies on Antibiotic Residues in Beef and Effect of Cooking and Freezing on Antibiotic Residues Beef Samples. *Scholarly Journal of Food and Nutritionm 2(1) 1-4*
66. 66.Shaltout FA, Zakaria IM and Nabil ME.( 2018): Incidence of Some Anaerobic Bacteria Isolated from Chicken

67. 67. Meat Products with Special Reference to Clostridium perfringens. *Nutrition and Food Toxicology* 2.5 (2018): 429-438.
68. 68. Luby S.P., Rahman M., Hossain M.J., Blum L.S., Husain M.M., Gurley E., Khan R., Ahmed B.N., Rahman S., Nahar N., Kenah E., Comer J.A., Ksiazek T.G. (2006): Foodborne transmission of Nipah virus, Bangladesh. *Emerg. Infect. Dis.*;12:1888–1894.
69. 69. Shaltout FA, Zakaria IM and Nabil ME. (2018): Incidence of Some Anaerobic Bacteria Isolated from Chicken Meat Products with Special Reference to Clostridium perfringens. *Nutrition and Food Toxicology* 2.5 (2018): 429-438.
70. 70. Shaltout FA, Ahmed A A Maarouf and Mahmoud ES Elkhouly. (2017): Bacteriological Evaluation of Frozen Sausage. *Nutrition and Food Toxicology* 1.5 ; 174-185.
71. 71. Shaltout FA, El-Toukhy EI and Abd El-Hai MM. (2019): Molecular Diagnosis of Salmonellae in Frozen Meat and Some Meat Products. *Nutrition and Food Technology Open Access* 5(1): dx.doi.org/10.16966/2470-6086.155.
72. 72. Shaltout, F.A., A.M. Ali and S.M. Rashad (2016): Bacterial Contamination of Fast Foods. *Benha Journal of Applied Sciences (BJAS)* 1 (2) 45-51
73. 73. Shaltout FA, Riad EM, TES Ahmed and Abou Elhassan Asmaa (2017): Studying the Effect of Gamma Irradiation on Bovine Offal's Infected with Mycobacterium tuberculosis Bovine Type. *Journal of Food Biotechnology Research* Vol.1 No.1:6: 1-5.
74. 74. Shaltout, F.A., Zakaria. I. M. , Jehan Eltanani , Asmaa . Elmelegy (2015): Microbiological status of meat and chicken received to University student hostel. *BENHA VETERINARY MEDICAL JOURNAL*, VOL. 29, NO. 2:187-192, DECEMBER, 201
75. 75. Saad, S.M.; Edris, A.M.; Shaltout, F.A. and Edris, Shima (2012): Isolation and identification of salmonellae and E. coli from meat and poultry cuts by using A. multiplex PCR. *Benha Vet. Med. J. special issue* 16-26
76. 76. Saad, S.M. and Shaltout, F.A. (1998): Mycological Evaluation of camel carcasses at Kalyobia Abattoirs  
a. *Vet. Med. J. Giza*, 46(3):223-229.
77. 77. Saad S.M. , Shaltout, F.A., Nahla A Abou Elroos, Saber B El-nahas. 2019: Antimicrobial Effect of Some Essential Oils on Some Pathogenic Bacteria in Minced Meat. *J Food Sci Nutr Res.* 2019; 2 (1): 012-020
78. 78. Saad S.M. , Hassanin, F. S; Shaltout, F.A., Marionette Z Nassif, Marwa Z Seif. (2019): Prevalence of Methicillin-Resistant Staphylococcus Aureus in Some Ready-to-Eat Meat Products. *American Journal of Biomedical Science & Research* 4(6):460-464.
79. 79. Saad S.M. , Shaltout, F.A., Nahla A Abou Elroos and Saber B El-nahas. (2019): Incidence of Staphylococci and E. coli in Meat and Some Meat Products. *EC Nutrition* 14.6.
80. 80. Lopman B.A., Reacher M.H., Van Duynhoven Y., Hanon F.X., Brown D., Koopmans M. (2003): Viral gastroenteritis outbreaks in Europe, 1995–2000. *Emerg. Infect. Dis.* ;9:90–96.
81. 81. Lupulović D., Lazić S., Prodanov-Radulović J., Jiménez de Oya N., Escribano-Romero E., Saiz J.C., Petrović T. (2010): First serological study of hepatitis E virus infection in backyard pigs from Serbia. *Food Environ. Virol.* ;2(2):110–113.
82. 82. Matsuda H., Okada K., Takahashi K., Mishiro S. (2003): Severe hepatitis E virus infection after ingestion of uncooked liver from a wild boar. *J. Infect. Dis.* ;188:944.
83. 83. Mattison K., Harlow J., Morton V., Cook A., Pollari F., Bidawid S., Farber J.M. (2010): Enteric viruses in ready-to-eat packaged leafy greens. *Emerg. Infect. Dis.* ;16(11):1815–1817.
84. 84. Mayr C., Strohe G., Contzen M. (2009): Detection of rotavirus in food associated with a gastroenteritis outbreak in a mother and child sanatorium. *Int. J. Food Microbiol.*;135(2):179–182.
85. 85. Shaltout FA, Mohamed, A. Hassan and Hassanin, F. S (2004): THERMAL INACTIVATION OF ENTEROHAEMORRHAGIC ESCHERICHIA COLI O157:H7 AND ITS SENSITIVITY TO NISIN AND LACTIC ACID CULTURES. 1st Ann. Confr. , FVM., Moshtohor, Sept, 2004
86. 86. Shaltout FA, El-diasty, E.M. ; Elmesalamy, M. and Elshaer, M. 2014: Study on fungal contamination of some chicken meat products with special reference to the use of PCR for its identification . Conference, *Veterinary Medical Journal – Giza* vol. December 2014/12/17 vol.60: 1-10
87. 87. Shaltout, F.A. 2002: Microbiological Aspects of Semi-cooked chicken Meat Products. *Benha Veterinary Medical Journal* 13,2,: 15-26
88. 88. Mesquita J.R., Vaza L., Cerqueira S., Castilho F., Santos R., Monteiro S., Manso C.F., Romalde J.L., Nascimento M.S.J. (2011): Norovirus, hepatitis A virus and enterovirus presence in shellfish from high quality harvesting areas in Portugal. *Food Microbiol.*;28:936–941
89. 89. Shaltout FA, Mohammed Farouk; Hosam A.A. Ibrahim and Mostafa E.M. Afifi. 2017: Incidence of Coliform and Staphylococcus aureus in ready to eat fast foods. *BENHA VETERINARY MEDICAL JOURNAL*, 32( 1): 13 - 17, MARCH, 2017
90. 90. Müller L., Schultz A.C., Fonager J., Jensen J., Lisby M., Hindsdal K., Krusell L., Eshøj A., Møller L.T., Porsbo L.J., Böttiger B.E., Kuhn K., Engberg J., Ethelberg S. (2014): Separate norovirus outbreaks linked to one source of imported frozen raspberries by molecular analysis, Denmark, 2010–2011. *Epidemiol. Infect.*;22:1–9.
91. 91. Shaltout, F.A. (1992): Studies on Mycotoxins in Meat and Meat by Products. M.V.Sc Thesis Faculty of Veterinary Medicine, Moshtohor, Zagazig University Benha branch.
92. 92. Shaltout, F.A. (1996 ): Mycological And Mycotoxicological profile Of Some Meat products. Ph.D. Thesis, Faculty of Veterinary Medicine, Moshtohor, Zagazig University Benha branch.
93. 93. Shaltout, F.A. (1998): Proteolytic Psychrotrophes in Some Meat products. *Alex. Vet. Med. J.* 14 (2):97-107.
94. 94. Shaltout, F.A. (1999): Anaerobic Bacteria in Vacuum Packed Meat Products. *Benha Vet. Med. J.* 10 (1):1-10
95. Shaltout, F.A. (2000): Protozoal Foodborne Pathogens in some Meat Products. *Assiut Vet. Med. J.* 42 (84):54-59.
96. 97. Shaltout, F.A. (2001): Quality evaluation of sheep carcasses slaughtered at Kalyobia abattoirs.
97. 98. Assiut Veterinary Medical Journal, 46(91):150-159.
98. 99. Shaltout, F.A. (2002): Microbiological Aspects of Semi-cooked Chicken Meat Products. *Benha Vet. Med. J.* 13(2):15-26.
99. 100. Shaltout, F.A. (2003): Yersinia Enterocolitica in some meat products and fish marketed at Benha city.
100. 101. The Third international conference Mansoura 29-30 April.
101. 102. Shaltout, F.A. (2009): Microbiological quality of chicken carcasses at modern Poultry plant
102. 103. The 3rd Scientific Conference, Faculty of Vet. Med. , Benha University, 1-3 January
103. 104. Newell D.G., Koopmans M., Verhoef L., Duizer E., Aidara-Kane A., Sprong H., Opsteegh M., Langelaar M., Threlfall J., Scheutz F., van der Giessen J., Kruse H. (2010): Food-borne diseases—the challenges of 20 years ago still persist while new ones continue to emerge. *Int. J. Food Microbiol.* ;139:S3–S15.

104. 105.Norder H., Sundqvist L., Magnusson L., Østergaard Breum S., Löfdahl M., Larsen L.E., Hjulsgaard C.K., Magnius L., Böttiger B.E., Widén F.( 2009) :Endemic hepatitis E in two Nordic countries. *Euro Surveill.*;14(19):1–9.
105. 106.Shaltout,F.A., Amin, R., Marionet , Z., Nassif and Shima, Abdel-wahab( 2014): Detection of aflatoxins in some meat products. *Benha veterinary medical journal* , 27( 2) :368-374
106. 107.Shaltout,F.A. and Afify , Jehan Riad,EM and Abo Elhasan , Asmaa,A.(2012): Improvement of microbiological status of oriental sausage. *Journal of Egyptian Veterinary Medical Association* 72(2):157-167
107. 108.Shaltout,F.A. and Daoud, J. R.(1996): Chemical analytical studies on rabbit meat and liver. *Benha Vet. Med.J.8* (2):17-27.
108. 109.Shaltout,F.A. and Edris, A.M.(1999): Contamination of shawerma with pathogenic yeasts. a. *Assiut Veterinary Medical Journal*,40(64):34-39.
109. 110.Shaltout, F. A. ;Eldiasty , E. and Mohamed , M.S.(2014): Incidence of lipolytic and proteolytic fungi in some chicken meat products and their public health significance. *Animal Health Research Institute : First International Conference on Food Safety and Technology 19-23 June 2014 Cairo Egypt* pages 79-89.
110. 111.Shaltout, F.A.;Eldiasty, E. ; Salem, R. and Hassan, Asmaa(2016): Mycological quality of chicken carcasses and extending shelf – life by using preservatives at refrigerated storage. *Veterinary Medical Journal -Giza (VMJG)*62(3)1-7
111. 112.Shaltout, F.A.; Salem, R. Eldiasty, E. ; and Diab, Fatema. (2016): Mycological evaluation of some ready to eat meat products with special reference to molecular characterization. *Veterinary Medical Journal -Giza* 62(3)9-14
112. 113.Shaltout, F. A. ;Elshater , M. and Wafaa , Abdelaziz (2015): Bacteriological assessment of street vended meat products sandwiches in Kalyobia Governorate. *Benha Vet. Med.J.28* (2):58-66.
113. 114.Shaltout, F. A. ; Gerges, M.T. and Shewail, A.A.(2014):Impact of Organic Acids and Their Salts on Microbial Quality and Shelf Life of Beef. *Global Journal of Agriculture and Food Safety Science* 1(2): 360-370. Massive publisher house M.P.H. Egypt [www.mphegypt.com](http://www.mphegypt.com)
114. 115.Shaltout,F.A.;Ghoneim, A.M.; Essmail, M.E. and Yousseif ,A.(2001): Studies on aflatoxin B1 residues in rabbits and their pathological effects. *J.Egypt. Vet. Med. Association* 61(2):85-103.
115. 116.Shaltout,F.A. and Hanan ,M.T. El-Lawendy (2003): Heavy Metal Residues In Shawerma.
116. 117.Beni-Suef Vet.Med.J. 13(1):213-224.
117. 118.Pebody R.G., Leino T., Ruutu P., Kinnunen L., Davidkin I., Nohynek H., Leinikki P. (1998):Food-borne outbreaks of hepatitis A in a low endemic country: an emerging problem? *Epidemiol. Infect.* ;120:55–59.
118. 119.Shaltout, F.A. and Hashim, M.F. (2002): Histamine in salted, Smoked and Canned Fish products. *Benha Vet. Med.J.13* (1):1-11.
119. 120.Shaltout, F.A. ; Hashim,M.F. and Elnahas,s.(2015): Levels of some heavy metals in fish (*tilapia nilotica* and *Claris lazera*) at Menufia Governorate. *Benha Vet. Med.J.29* (1):56-64
120. 121.Shaltout,F.A. and Ibrahim, H.M.(1997): Quality evaluation of luncheon and Alexandrian sausage. *Benha Vet. Med.J.10* (1):1-10.
121. 122.Shaltout, F.A. ; Nassif, M and Shakran , A(2014): Quality of battered and breaded chicken meat products
122. 123.*Global Journal of Agriculture and Food Safety Science – 1* ( 2 ) ISSN 2356-7775
123. 124.Shaltout,F.A., Amani M. Salem, A. H. Mahmoud, K. A(2013): Bacterial aspect of cooked meat and offal at street vendors level. *Benha veterinary medical journal*, 24(1): 320-328
124. 125.Shaltout,F.A. and Salem, R.M.(2000):Moulds, aflatoxin B1 and Ochratoxin A in Frozen Livers and meat products. *Vet . Med. J.Giza* 48(3):341-346.
125. 126.Tamada Y., Yano K., Yatsuhashi H., Inoue O., Mawatari F., Ishibashi H. (2004):Consumption of wild boar linked to cases of hepatitis E. *J. Hepatol.* ;40:869–870.
126. 127.Yasser H. Al-Tarazi, A. Al-Zamil, Shaltout FA. and H. Abdel- Samei (2002). Microbiological status of raw cow milk marketed in northern Jordan. The sixth Scientific Conference of Zagazig University, Faculty of veterinary Medicine, held at AlQarah- Egypt, 7-9 Sept.
127. 128.Petrignani M., Harms M., Verhoef L., van Hunen R., Swaan C., van Steenberg J., Boxman I., Peran i Sala R., Ober H.J., Vennema H., Koopmans M., van Pelt W. Update (2010): a food-borne outbreak of hepatitis A in the Netherlands related to semi-dried tomatoes in oil, January-February 2010. *Euro Surveill.* ;15(20).
128. 129.Shaltout, F. A.; El-diasty, E.M. and Mohamed, M. S.(2014): Incidence of lipolytic and proteolytic fungi in some chicken meat products and their public health significance. *1st Scientific conference of food safety and Technology .2014*, pp. 79-89
129. 130.Petrović T., Lupulović D., Jimenez de Oya N., Vojvodić S., Blazquez A., Escribano-Romero E., Martin-Acebes M., Potkonjak A., Milošević V., Lazić S., Saiz J.C. (2014):Prevalence of hepatitis E virus antibodies in Serbian blood donors. *J. Infect. Dev. Ctries.* ;8(10):1322–1327.
130. 131.Shaltout FA, R.M. Salem, E.M. El-Diasty and W.I.M. Hassan. 2019: Effect of Lemon Fruits and Turmeric Extracts on Fungal Pathogens in Refrigerated Chicken Fillet Meat. *Global Veterinaria* 21 (3): 156-160,
131. 132.Richards G.P. (1985):Outbreaks of shellfish—associated enteric virus illness in the United States: requisite for development of viral guidelines. *J. Food Prot.* ;48:815–823.
132. 133.Ramsay C.N., Upton P.A. (1989):Hepatitis A and frozen raspberries. *Lancet.* ;1:43–44.
133. 134.Wheeler C., Vogt T.M., Armstrong G.L., Vaughan G., Weltman A., Nainan O.V., Dato V., Xia G., Waller K., Amon J., Lee T.M., Highbaugh-Battle A., Hembree C., Evenson S., Ruta M.A., Williams I.T., Fiore A.E., Bell B.P. ( 2005):An outbreak of hepatitis A associated with green onions. *N. Engl. J. Med.* ;353:890–897.
134. 135.Shaltout FA, Ahmed, A.A. Maarouf , Eman, M.K. Ahmed(2018): Heavy Metal Residues in chicken cuts up and processed chicken meat products. *BENHA VETERINARY MEDICAL JOURNAL*, 34( 1): 473-483.
135. 136.Shaltout ,F.A.; Hanan M. Lamada , Ehsan A.M. Edris.(2020): Bacteriological examination of some ready to eat meat and chicken meals. *Biomed J Sci & Tech Res.*, 27(1): 20461- 20465
136. 137.Sobhy, Asmaa and Shaltout, Fahim(2020): Prevalence of some food poisoning bacteria in semi cooked chicken meat products at Qaliubiya governorate by recent Vitek 2 compact and PCR techniques. *Benha Veterinary Medical Journal* 38 (2020) 88-92.
137. 138.Sobhy, Asmaa and Shaltout, Fahim(2020): Detection of food poisoning bacteria in some semi-cooked chicken meat products marketed at Qaliubiya governorate. *Benha Veterinary Medical Journal* 38 (2020) 93-96
138. 139.Holzmann H., Aberle S. W., Stiasny K., Werner P., Mischak A., Zainer B., Heinz F. X. (2009) : Tick-borne encephalitis from eating goat cheese in a mountain region of Austria. *Emerg Infect Dis.*;15.10:1671.

139. 140. Cisak E, Wójcik-Fatla A, Zajac V, Sroka J, Buczek A, Dutkiewicz J. (2010): Prevalence of tick-borne encephalitis virus (TBEV) in samples of raw milk taken randomly from cows, goats and sheep in eastern Poland. *Ann Agric Environ Med*;17(2):283-86.
140. 141. Koopmans M, Duizer E. (2004): Foodborne viruses: an emerging problem. *Int J Food Microbiol*;90(1):23-41.
141. 142. Vasickova P, Psikal I, Kralik P, Widen F, Hubalek Z, Pavlik I. (2007) Hepatitis E virus: a review. *Vet Med-Czech*;52:365-84.
142. 143. Todd ECD, Greig JD, Bartleson CA, Michaels BS. (2007) Outbreaks where food workers have been implicated in the spread of foodborne disease. Part 2. description of outbreaks by size, severity, and settings. *J Food Prot.*;70:1975-93.
143. 144. Todd ECD, Greig JD, Bartleson CA, Michaels BS. (2007) Outbreaks where food workers have been implicated in the spread of foodborne disease. Part 3. factors contributing to outbreaks and description of outbreak categories. *J Food Prot.*;70:2199-2217.
144. 145. Ranst M, Robesyn E, Schrijver KD, Top G, Verbeeck J, Wollants E. (2009): An Outbreak of Hepatitis A Associated with the Consumption of Raw. Beef. *J Clin Virol*;44(3):207-10.
145. 146. Mattison K, Shukla A, Cook A, Pollari F, Friendship R, Kelton D, Bidawid S, Farber JM. (2007): public noroviruses in swine and cattle. *Emerg Infect. Dis*;13:1184-88.
146. 147. Malek M et al. (2009) Outbreak of norovirus infection among river rafters associated with packaged delicatessen meat, Grand Canyon, 2005. *Clin. Infect Dis*;48(1):31-7.
147. 148. Nordgren J, Kindberg E, Lindgren P-E, Matussek A, Svensson L. Norovirus (2010): Gastroenteritis Outbreak with a Secretor-independent. Susceptibility Pattern, Sweden. *Emerg Infect Dis*;16(1):81-87.
148. 149. Radin D, Velebit B. (2015): Transmission of foodborne viruses during food handling. In Press Invited paper. *Microbiologia Balkanica*, Greece.
149. 150. Kingsley DH, Chen H. (2009) Influence of pH, salt, and temperature on pressure inactivation of hepatitis A virus. *Int J Food Microbiol*;130:61-4.
150. 151. Hudson JB, Sharma M, Vimalanathan S. (2009): Development of a practical method for using ozone gas as a virus decontaminating agent. *Ozone Sci. Eng*;31:216-23.
151. 152. Feagins AR, Opriessnig T, Guenette DK, Halbur PG, Meng XJ. (2008): Inactivation of infectious hepatitis E virus present in commercial pig livers sold in local grocery stores in the United States. *Int J Food Microbiol*;123:32-7.



This work is licensed under Creative Commons Attribution 4.0 License

To Submit Your Article Click Here:

**Submit Manuscript**

DOI: [10.31579/2642-9756/163](https://doi.org/10.31579/2642-9756/163)

#### Ready to submit your research? Choose Auctores and benefit from:

- fast, convenient online submission
- rigorous peer review by experienced research in your field
- rapid publication on acceptance
- authors retain copyrights
- unique DOI for all articles
- immediate, unrestricted online access

At Auctores, research is always in progress.

Learn more <https://www.auctoresonline.org/journals/women-health-care-and-issues>