

Effect of dietary supplementation of *Monotheca buxifolia* leaf powder on the growth performance of broiler chickens

Alagbe Olujimi John

Department of Animal Nutrition and Biochemistry, Sumitra Research Institute, Gujarat, India

Corresponding author: Alagbe Olujimi John, Department of Animal Nutrition and Biochemistry, Sumitra Research Institute, Gujarat, India.

Received date: August 16, 2024; **Accepted date:** September 04, 2024; **Published date:** September 20, 2024

Citation: Alagbe O. John, (2024), Effect of dietary supplementation of *Monotheca buxifolia* leaf powder on the growth performance of broiler chickens, *J. Nutrition and Food Processing*, 7(11); DOI:10.31579/2637-8914/258

Copyright: © 2024, Alagbe Olujimi John. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract:

The aim of this study was to investigate the effect of dietary supplementation of *Monotheca buxifolia* leaf powder on the growth performance of broiler chickens. A total of 250 -1-day old broiler chicks (Ross 308) were randomly distributed into five treatments with five replicates and each replicate consists of ten birds. Prior to the arrival of birds, basal diet adequate in all nutrients was formulated according to the requirement of birds by Nutritional Research Council in 1994. Birds in treatment one was fed basal diet without *Monotheca buxifolia* leaf powder, those in treatment two, three, four and five were fed same diet with *Monotheca buxifolia* leaf powder at 100 g, 200 g, 300 g and 400 g respectively. The experiment lasted for 42 days and birds had unrestricted access to clean water and feed. A completely randomized experimental design was adopted and all management practices was strictly adhered to. Result on the phytochemical content of *Monotheca buxifolia* leaf powder flavonoids (863.2 mg/g), phenols (511.8 mg/g) and terpenoids (102.3 mg/g) were the most prominent compounds followed by saponins (72.56 mg/g), tannins (62.34 mg/g), alkaloids (41.88 mg/g) and steroids (30.41 mg/g). Average daily weight gain and average daily feed intake values which varied from 41.24 – 51.37 g/b and 103.7 – 107.5 g/b were higher among birds fed *Monotheca buxifolia* leaf powder (treatment two, three, four and five) relative to treatment one ($P < 0.05$). Similarly, feed conversion ratio whose value ranged from 2.09 – 2.50 were significantly ($P < 0.05$) influenced by the treatment. In conclusion, feeding broilers *Monotheca buxifolia* leaf powder up to 400 g/kg diet does not pose any detrimental effect on the performance of birds.

Key words: *Monotheca buxifolia*; phytochemicals; growth; performance; food safety; dose

Introduction

Using antibiotics in sub therapeutic dosages was for a long time a general tool for the control of diseases, however, their misuse in animal husbandry has contributed to the development of multiple resistance of pathogens, deposit of drug residues in edible animal product and the environment, and the potential transfer of antibiotic resistance to human pathogens has directed research towards alternative solutions such as the use of medicinal or herbal plants (Daniel, 2020; Peter, 2021). Plants are natural reservoir of medicinal agents almost free from the side effects normally caused by synthetic chemicals (Fennel et al., 2004). They also contain a wide variety of free radicals scavenging molecules including phenols, flavonoids, vitamins, terpenoids that are rich in antioxidant activity (Cai and Sun, 2003). Medicinal plants are known to contain phytochemicals which can be found in stems, roots, leaves, stem bark, flowers amongst others, possess numerous pharmacological properties, antimicrobial, anti-fungal, antioxidant, gastro-protective, cytotoxic, hypolipidemic, anti-nociceptive, antidiuretic, immune-stimulatory, anti-inflammatory and antiviral (Alagbe, 2023; John, 2024).

Monotheca buxifolia is one of the numerous underexplored medicinal plant belonging to the family Sapotaceae (Maryam et al., 2020). The plant

is widely distributed in Pakistan, Afghanistan, Oman, Saudi Arabia and some parts of India (Ihsan et al., 2020). Extracts from the leaves, stem and roots of the plant can be used for the traditional treatment of digestive disorders, urinary tract disease, fever, sexually transmitted infections, diabetes, peptic ulcers, piles, yaws, dysmenorrhea, infertility, and helminthic infections (Irfan et al., 2016; Jan et al., 2013). Uses of its root, root bark and bark of stem are extensive, particularly for their astringent, haemostatic, hypotensive, vulnerary and diaphoretic activities (Rehman et al., 2013). The leaves contain, flavonoids, terpenoids, glycosides, phenolic compound, tannins and anthraquinones at different concentrations which contributes to their several therapeutic or biological functions, anti-inflammatory, antioxidant, cytotoxic, phytotoxic, anti-pyretic, central nervous system depressant, and hepato-protective (Ullah et al., 2016). This is possible because concentration of plant constituents of the same plant organ can vary from one geographical location to another depending on the age of the plant, differences in topographical factors, the nutrient concentrations of the soil, extraction method (Alagbe, 2023; Ojediran et al., 2024). Ethanolic and methanolic extracts from the leaves of *Monotheca buxifolia* possesses antimicrobial properties and

have been reported to inhibit the activities of *Escherichia coli*, *Klebsiella* spp, *Salmonella* spp and *Staphylococcus* spp (Ullah et al., 2016).

Previous studies by notable researchers have shown that phytochemicals exerts positive influence on the growth performance, immune response and microbial population in the gastrointestinal tract of birds (Oloruntola et al., 2016; Kanduri et al., 2013; Hashemi and Davoodi, 2010; Rabelo et al., 2003; Seidavi and Simoes, 2015). However, outcome of their findings have not been consistent, this could be due to differences in inclusion levels, specie of plant used as well as their chemical constituents (Adewale et al., 2021). There is little or no report on the effect of *Monothecha buxifolia* leaf powder on the growth performance of broiler chickens. This research is timely as it will help to address the increasing cases of antimicrobial resistance, provide optimum levels for birds and help to promote food safety.

Materials and methods

Experimental area

The experiment was carried out at the Poultry Section, Sumitra Research Institute, Gujarat, India between the month of January to March, 2023. The institute is located between 23° 13' N and 72° 41' E.

Collection and preparation of *Monothecha buxifolia* leaf powder

Freshly harvested leaves of *Monothecha buxifolia* were harvested from Orathur village in Kancheepuram district, India. The collected leaves were washed with running tap water and air dried in an open shade for 10 days after authentication at the department of taxonomy, Sumitra Research Institute, Gujarat and assigned a reference number (GB/056F/2023). The dried leaves were grounded into powder with electric blender and packed into a labeled polythene bag before it was sent to the laboratory for further analysis.

Management of experimental animal and design

This study was carried out according to the guidelines of animal protocol approved by the Research and Ethics Committee of the department of Animal Nutrition and Biochemistry, Sumitra Research Institute, Gujarat,

India. 250 – 1-day old broiler chick (Ross 308) mixed sex with an average initial body weight of 51.2 ± 0.02 g randomly distributed to five treatments with five replicates consisting of 10 birds each. Birds were reared in a battery cages equipped with nipple drinkers and manual feeders kept in a semi-closed pens. Prior to the arrival of bird's, battery cages were properly disinfected and a basal diet which is adequate in all nutrient was formulated according to Nutritional Research Council's guidelines (1994) for broilers. A completely randomized experimental design was adopted with birds in treatment 1 fed basal diet without *Monothecha buxifolia* leaf powder while those in treatment 2, 3, 4 and 5 were fed same diet with *Monothecha buxifolia* leaf powder at 100 g, 200 g, 300 g and 400 g/kg diet. Birds had unrestricted access to clean water and feed. Weight gain and feed intake were taken into consideration throughout the experiment which lasted for 42 days.

Determination of phytochemical contents in *Monothecha buxifolia* leaf powder

Quantification of flavonoids, alkaloids, saponins, steroids, tannins and terpenoids contents in *Monothecha buxifolia* leaf powder was carried out according to procedures recently published by Alagbe (2024). Each phyto-constituents were recorded at different optical densities using GC/MS Tripod (Model 821W-011J, China).

Proximate content of experimental diet

Proximate content of experimental diet was carried out using near infrared automated kit (NIR -7000, USA) which uses SensorVu windows® based PC software. All operations were carried out according to the manufacturers recommendation.

Statistical analysis

All the data obtained were subjected to one-way analysis of variance (ANOVA) using SPSS version 25. The differences among the treatment means were determined ($P < 0.05$) by Duncan multiple range test of the same software.

Results and discussion

Ingredients	Starter mash (0-28 d)	Finishers mash (29 – 42 d)
Corn	50.00	52.00
Wheat bran	5.60	8.00
Soya bean	34.00	30.00
Fish meal	6.20	5.00
Di-calcium phosphate	2.50	3.00
Limestone	1.20	1.50
DL-methionine	0.25	0.20
Lysine	0.25	0.20
^{1,2} *Mineral-vitamin premix	0.25	0.25
Salt	0.30	0.35
Total	100.0	100.0
Chemical composition		
Dry matter	91.29	91.04
Crude protein	23.04	21.22
Crude fibre	4.20	5.06
Ether extract	3.92	3.55
Methionine	0.50	0.62
Methionine +cysteine	0.92	0.98
Lysine	1.17	1.19
Ash	6.92	7.11
Calcium	1.43	1.61
Phosphorus	0.61	0.72
Metabolizable energy (kcal/kg)	2907.1	3204.8

¹2.5 kg Mineral/vitamin premix for starter contains: Vit. A, 12000000 IU; Vit.D3 1800000 IU; Vit.E, 15000 mg; Vit.K3, 1000 mg; Vit.B1, 1200 mg; Vit.B2, 5100 mg; Vit. B6, 1500 mg; Vit.B12, 10mg; biotin, 50mg; pantothenic acid, 10000 mg; nicotinic acid, 30000 mg; folic acid, 1000 mg; choline chloride, 250000 mg; Mn, 60000 mg; Zn, 50000 mg; Fe, 30000 mg; Cu, 10000 mg; I, 1000 mg; Se, 100mg; Co, 100mg

²2.5 kg Mineral/vitamin premix for finisher contains: Vit. A, 12000000 IU; Vit.D3 2000000 IU; Vit.E, 10000 mg; Vit.K3, 2000 mg; Vit.B1, 1000 mg; Vit.B2, 5000 mg; Vit. B6, 1500 mg; Vit.B12, 10mg; biotin, 50mg; pantothenic acid, 10000 mg; nicotinic acid, 30000 mg; folic acid, 1000 mg; choline chloride, 250000 mg; Mn, 60000 mg; Zn, 50000 mg; Fe, 30000 mg; Cu, 10000 mg; I, 1000 mg; Se, 100mg; Co, 100 mg

Table 1: Ingredient and chemical composition of experimental diet (as fed basis)

Phytochemical content of *Monotheca buxifolia* leaf powder is presented in Table 2. Flavonoids had the highest concentration of 863.2 mg/g followed by phenols (511.8 mg/g), terpenoids (102.3 mg/g), saponins (72.56 mg/g), tannins (62.34 mg/g), alkaloids (41.88 mg/g) and steroids (30.41 mg/g). The presence of these phyto-constituents showed that *Monotheca buxifolia* leaf possesses several medicinal or pharmacological properties (Singh et al., 2022; John, 2024). Daniel et al. (2024); Adewale et al. (2021) reported that the concentration of phyto-constituents in medicinal plants are influenced by age of plant, geographical location, species, processing methods amongst others. Concentration of flavonoids and phenolic compound recorded in this study was higher than those reported for *Dysphania ambrosioides* (2.29 mg/g, 15.24 mg/g) and *Crassocephalum crepidioides* leaves (1.62 mg/g, 13.07 mg/g) by Falowo et al. (2023). Phenols and flavonoids have been suggested to have antioxidant properties (Dhan et al., 2007) and these antioxidants exert their activity by scavenging the „free oxygen radicals“ thereby giving rise

to a fairly „stable radical“. The free radicals are metastable chemical species, which tend to trap electrons from the molecules in the immediate surroundings. These radicals if not scavenged effectively in time, they may damage crucial bio molecules like lipids, proteins including those present in all membranes, mitochondria and, the DNA resulting in abnormalities leading to disease conditions (Uddin et al. 2008). Other properties include; antimicrobial (Dandan, 2009), anticarcinogenesis (Aritra and Sumana, 2012), anti-inflammatory (Daljit and Gurinder, 2007), cardio-protective (Alagbe, 2024) and immune-stimulatory (Gupta et al., 2003). Saponins and terpenoids have been suggested to possess hypolipidemic and anticancer properties (Kris-Etherton et al., 2002). Alkaloids have pharmacological applications as analgesics, antimalarial and central nervous system stimulants (Madziga et al., 2010; John, 2024). The result obtained in this study is in agreement with the reports of Maryam et al. (2020); Ullah et al. (2016).




Table 2: Phytochemical content of *Monotheca buxifolia* leaf powder

Effect of *Monotheca buxifolia* leaf powder on the growth performance of broiler chicken is presented in Table 3. Average daily weight gain, average daily feed intake and feed conversion ratio were significantly ($P < 0.05$) influenced by the treatments. Birds fed *Monotheca buxifolia* leaf powder had higher values compared to the control in T1 ($P < 0.05$). Results obtained suggests that *Monotheca buxifolia* leaf powder was able to modulate the activities of digestive enzymes thus leading to efficient nutrient utilization among birds. This action is triggered by the presence of phytochemicals in the test ingredient (*Monotheca buxifolia* leaf) (Omokore and Alagbe, 2019; Agubosi et al., 2022). Outcome also suggests that *Monotheca buxifolia* leaf possess antibacterial properties thus preventing dysbiosis which could also translate to a better feed conversion ratio in birds (Musa et al., 2021). The average daily weight

gain range observed in this study with the dietary supplementation of *Monotheca buxifolia* leaf powder 41.24 – 51.37 g/b was similar to the result of a study by John (2024); Alagbe (2019) who discovered that average daily weight gains of broilers fed *Rhamnus prinoides* leaf extract varied from 40.08 – 55.71 g/b. The result was higher than those presented by Agubosi et al. (2021) when sunflower oil was supplemented in the diet of broiler chicken at 0.3 %. Findings of this study on average daily feed intake (103.1 – 107.5 g/b) were lower than 81.18 – 82.30 g/b reported by Oloruntola et al. (2021); Kholoud et al. (2021) when phyto-additives were fed to broiler chickens but similar to 99.60 – 114.8 g/b recorded by Al-Mufarrej et al. (2019) when clove powder was supplemented in the diet of broilers. Feed conversion ratio values which varied from 2.09 – 2.50 was similar to the results of a study by Dibaji et al. (2014) who found that feed conversion ratio of broilers fed symbiotic ranged from 2.00 – 2.20.

This result was lower than those reported by Goliomytis et al. (2014) when quercetin was fed to broiler chickens.

Variables	T1	T2	T3	T4	T5	SEM
Number of birds per treatment	50.00	50.00	50.00	50.00	50.00	-
Duration of experiment (days)	42.00	42.00	42.00	42.00	42.00	-
Initial body weight (g/bird)	51.2	51.22	51.03	51	51.01	0.01
Final body weight (g/bird)	1783.1	2206.7	2207.5	2208.3	2208.5	61.82
¹ Weight gain (g/bird)	1731.9	2155.4	2156.5	2157.3	2157.5	58.07
² Average daily weight gain (g/bird)	41.24	51.33	51.34	51.36	51.37	0.02
³ Total feed intake (g/bird)	4328.7	4506.7	4511.2	4513.2	4515.1	91.14
⁴ Average daily feed intake (g/bird)	103.1	107.3	107.4	107.5	107.5	0.05
⁵ Feed conversion ratio	2.50	2.09	2.09	2.09	2.09	0.01

Means on the same row having different superscripts are significantly different ($P < 0.05$); SEM: standard error of mean; T1: basal diet without *Monotheca buxifolia* leaf powder; T2, T3, T4 and T5: basal diet supplemented with *Monotheca buxifolia* leaf powder at 100 g, 200 g, 300 g and 400 g per kg diet respectively. ¹Final body weight – initial body weight ;²Average daily weight gain /42 days ;³Feed served – left over; ⁴Total feed intake/42 days; ⁵Average daily feed intake/average daily weight gain.

Table 3: effect of *Monotheca buxifolia* leaf powder on the growth performance of broiler chicken

Conclusion

In conclusion, *Monotheca buxifolia* leaf powder contains several phytoconstituents with medicinal value with flavonoids, phenols and terpenoids dominating as major bioactive compounds. These compounds perform multiple biological activities such as, anti-inflammatory, anti-pyretic, antioxidant, antifungal, cytotoxic, gastro-protective, immuno-stimulatory functions amongst others. Dietary supplementation of *Monotheca buxifolia* leaf powder to broilers up to 400 g/kg diet had no negative effect on their growth performance.

References

- Kholoud, O.A., Intisar, Y.K and Mohammed, A. (2021). Effect of supplemented graded levels of clove extracts on broiler chick's performance. *Journal of Animal Sciences and Livestock Production*, 5(4): 8812.
- Kuete V, Tangmouo JG, Penlap Beng V, Ngounou FN, Lontsi D. (2006). Antimicrobial activity of the methanolic extract from the stem bark of *tridesmostemon omphalocarpoides* (Sapotaceae). *Journal of Ethnopharmacology*, 104(1–2):5–11.
- Ullah I, Khan JA, Adhikari A, Khan A, Hannan PA, Wadood I. (2006). Bioassay guided isolation of new urease inhibitory constituents from *Monotheca buxifolia* (Falc.) fruit and their molecular docking studies. *Animal Production*, 10(6):744–749.
- Ullah I, Khan JA, Adhikari A, Shahid M. (2016). Hepatoprotective effect of *Monotheca buxifolia* fruit against antitubercular drugs-induced hepatotoxicity in rats. *Bangladesh Journal of Pharmacology*, 11(1):248–256.
- Jan S, Khan MR, Rashid U, Bokhari J. (2013). Assessment of antioxidant potential, total phenolics and flavonoids of different solvent fractions of *Monotheca Buxifolia* fruit. *Osong Public Health Research and Perspective*, 4(5):246–254.
- Rehman J, Khan IU, Farid S, Kamal S, Aslam N. (2013). Phytochemical screening and evaluation of in-vitro antioxidant potential of *Monotheca buxifolia*. *E3 Journal of Biotechnology and Pharmaceutical Research*, 4(4):54–60.
- Daniel, M. (2020). Synergistic effect of bioactive herbal extracts in gut stabilization. *International Poultry Magazine*, 2(3): 5-7.
- Peter, K. (2021). In-feed probiotic versus sub-therapeutic antibiotics. *International Pig Magazine*, 5(2): 3-4.
- Irfan, U., Jamshaid, A.K., Muhammad, S., Ajmal, K., Achyut, A., Peer, A.H, Ibrahim, J., Faisal, S and Umar, K. (2016). Pharmacological screening of *Monotheca buxifolia* (Falc.) A. DC. for antinociceptive, anti-inflammatory and antipyretic activities. *BMC Complementary and Alternative Medicine* (2016) 16:273.
- Ihsan, K., Joham, S.A and Ihsanm, U. (2020). Biological and phytochemical properties of *Monotheca buxifolia*: an unexplored medicinal plant, *Pharmaceutical Chemistry Journal*
- Oloruntola, O.D., Adu, O.A., Gbore, F.A., Falowo, A.B and Olarotimi, O.J. (2021). Performance of broiler chicken fed diets supplemented with *Irvingia gabonensis* kernel powder and *Ocimum gratissimum* leaf powder. *Slovakian Journal of Animal Science*, 54, 2021 (1): 7–20
- Al-Mufarrej, S.I., Fazea, E.H., Al-Baadani, H.H and Qaid, M.M. (2019). Effect of clove powder supplementation on performance, blood chemistry, and immune responses in broiler chickens. *South Africa Journal of Animal Science*, 49(5): 836-844.
- Dibaji, S. M., Seidavi, A., Asadpour, L. & Mareira da Silva, F. (2014). Effect of a symbiotic on the intestinal microflora of chickens. *Journal of Applied Poultry Research*, 23, 1–6.
- Goliomytis, M., Tsourekis, D., Simitzis, P. E., Charismiadou, M. A., Hager-Theodorides, A. L. & Deligeorgis, S. G. (2014). The effects of quercetin dietary supplementation on broiler growth performance, meat quality, and oxidative stability. *Poultry Science*, 93, 1957–1962.
- Madziga, H.A., Sanni, S. & Sandabe, U.K. (2010). Phytochemical and Elemental Analysis of *Acalypha wilkesiana* Leaf. *Journal of American Science*. 6(11): 510-514.

16. Uddin, S.N., Akond, M.A., Mubassara, S. & Yesmin, M.N. (2008). Antioxidant and Antibacterial activities of *Trema cannabina*. Middle-East Journal of Scientific Research. 3: 105-108.
17. Aritra Chatterjee and Sumana Chatterjee (2012). proximate analysis, phyto-chemical screening and anti-inflammatory activity of *coccinia indica*, International Journal of Pharmaceutical Chemical Biological Sciences. 2(3), 299-304.
18. Daljit, S.A. & Gurinder, J.K. (2007). Antibacterial activity of some Indian medicinal plants. Journal of Natural Medicines. 61(3):313-317.
19. Dandan L. (2009). Chemical composition and antimicrobial activity of essential oil isolated from the cultured mycelia of *Ganoderma japonicum*. Journal of Nanjing Medical University. 23(3): 168-172.
20. Dhan Prakash, Samiksha Suri, Garima Up adhyay & Singh, B.N. (2007). Total phenol, antioxidant and free radical scavenging activities of some medicinal plants. International Journal of food Sciences and Nutrition. 58(1): 18-28.
21. Gupta, M., Mazumder, U.K., Kumar, R.S. & Kumar, T.S. (2003). Studies on anti-inflammatory, analgesic and antipyretic properties of methanol extract of *Caesalpinia bonducella* leaves. Iranian Journal of Pharmacology and Therapeutics. 2: 30-33.
22. Kris-Etherton, P, M., Hecker, K.D., Bonanome, A., Coval, S.M., Binkoski, A.E., Hilpert, K.F., Griel, A.E. & Etherton, T.D. (2002). Bioactive compounds in foods: their role in the prevention of cardiovascular disease and cancer. American Journal of Medicine. 113:71S-88S.
23. Falowo, A., Oloruntola, O.D and Akinmoladun, O.F. (2023). Assessment of the nutritional composition and antioxidant properties of *Dysphania ambrosioides* and *Crassocephalum crepidioides* leaf as feed additives. Turkish Journal of Agriculture, 11(2): 274-279.
24. John, A.O., Muritala, S.D., Aduragbemi, Y.A., Kadiri, C.M., Bamiloye, S.O, Jummai, C and Effiong, E. (2024). The approximate mineral and phytochemical content of the leaves, stem bark and roots of *Pterocarpus erinaceus* in India. Cerrado: Agricultural and Biological Research, 1(1), 32-41.
25. John, A.O. (2024). Growth performance, haemato-biochemical indices of broiler chicken fed *Aristolochia indica* as a phyto-genic feed additive. Cerrado: Agricultural and Biological Research, 1(1), 42-53.
26. John, A.O. (2024). *Clerodendron splendens* leaf extract supplementation in weaner rabbits: impact on growth performance, haematology and intestinal microbial population. Cerrado: Agricultural and Biological Research, 1(1), 21-31.
27. John, A.O. (2024). Effect of coconut shell extract on the growth performance and some haemato-biochemical parameters of broiler chicken. Brazilian Journal of Science, 3(6): 82-95.
28. John, A.O. (2024). Impact of dietary supplementation of *Rhamnus prinoides* leaf extract on the growth performance, nutrient retention and intestinal microbial count of Japanese quails. Brazilian Journal of Science, 3(5), 40-50.
29. John, A.O. (2024). Effect of performance, serum biochemistry and haematological components of feeding Japanese quails phyto-genic feed additions comparing *Megaphrynium macrostachyum* leaves. Brazilian Journal of Science, 3(5), 51-64.
30. Musa, B., Alagbe, J.O., Adegbite Motunrade Betty & Omokore, E.A. (2020). Growth performance, caeca microbial population and immune response of broiler chicks fed aqueous extract of *Balanites aegyptiaca* and *Alchornea cordifolia* stem bark mixture. United Journal for Research and Technology, 2(2), 13-21.
31. Singh Sharma., Alagbe Olujimi John., Liu Xing., Sharma Ram and Kumar Amita (2022). Comparative analysis of ethanolic *Juniperus thurifera* leaf, stem bark and root extract using gas chromatography and mass spectroemetry. International Journal of Agriculture and Animal Production, 2(6), 18-27.
32. Shittu, M.D., Alagbe, J.O., Adejumo, D.O., Ademola, S.G., Abiola, A.O., Samson, B.O and Ushie, F.T. (2021). Productive Performance, Caeca Microbial Population and Immune-Modulatory Activity of Broiler Chicks Fed Different Levels *Sida Acuta* Leaf Extract in Replacement of Antibiotics. Bioinformatics and Proteomics Open Access Journal 5(1), 000143.
33. Ojediran, T.K., Emiola, I.A., Durojaye, V and John Alagbe (2024). Proximate, vitamin and GC-MS profiling of *Kigella africana* powder. Cerrado: Agricultural and Biological Research, 1(1): 13-20.
34. Ojediran, T.K., Emiola, I.A., Durojaye, V and John Alagbe (2024). Analysis of *Kigellia africana* fruit's powder antioxidant and phytochemical properties. Brazilian Journal of Science, 3(7): 38-49.
35. Agubosi, O.C.P., Alexander, James and Alagbe, J.O. (2022). Influence of dietary inclusion of Sunflower (*Helianthus annus*) oil on growth performance and oxidative status of broiler chicks. Central Asian Journal of Medical and Natural Sciences, 2(7), 187-195.
36. Agubosi, O.C.P., Soliu, M.B and Alagbe, J.O. (2022). Effect of dietary inclusion levels of *Moringa oleifera* oil on the growth performance and nutrient retention of broiler starter chicks. Central Asian Journal of Theoretical and Applied Sciences, 3(3), 30-39.
37. Adewale, A.O., Alagbe, J.O & Adeoye, Adekemi. O. (2021). Dietary Supplementation of *Rauvolfia Vomitoria* Root Extract as A Phyto-genic Feed Additive in Growing Rabbit Diets: Haematology and serum biochemical indices. International Journal of Orange Technologies, 3(3), 1-12.
38. Alagbe Olujimi John (2024). Novel phyto-genics' impact on weaned pigs growth performance, haematology and serum biochemical indicators. Black Sea Journal of Agriculture, 7(2), 82-89.
39. Alagbe, J.O., Shittu, M.D & Tanimomo, Babatunde K. (2022). Influence of *Anogeissusleio carpus* stem bark on the fatty acid composition in meat of broiler chickens. European Journal of Life Safety and Stability, 14(22), 13-22.
40. Daniel Nnadozie Anorue, Friday Ubong & Alagbe Olujimi John (2023). Investigating the effects of pawpaw (*Carica papaya*) essential oil dietary supplementation on the growth performance and carcass characteristics of broilers. Research in: Agricultural and Veterinary Sciences, 7(3), 164 - 174.

41. Cai, Y.Z and Sun, M. (2003). Antioxidant activity of betalins from plants of the Amaranthacea. *Journal of Agriculture and Food Chemistry*. 51: 2288-2294.
42. Hashemi, S. R and Davoodi, H. (2010). Phytochemicals as new class of feed additive in poultry industry. *Journal of Animal and Veterinary Advances*, 9, 2295–9304.
43. Rabelo, M., Souza, E. P., Soares, P. M., Miranda, A. V., Matos, F. J and Criddle, D. N (2003). Antinociceptive properties of the essential oil of *Ocimum gratissimum* L. (Labiatae) in mice. *Brazilian Journal of Medical and Biological Research*, 36, 521–524.
44. Seidavi, A and Simoes, J. (2015). Evaluation of dietary fish oil plus green tea supplementation on the gizzard, ileum and cecum microflora in broiler chickens. *Archivos de Zootecnia*, 64, 397–402.
45. Alagbe, J.O., Shittu, M.D., Ramalan, S.N., Tanimomo, K.B and Adekunle, D.A. (2022). Growth performance, semen quality characteristics and hormonal profile of male rabbit bucks fed *Rubia cordifolia* root extracts. *International Journal of Biological Engineering and Agriculture* 1(1): 1-13.
46. Omokore, E.O and Alagbe, J.O. (2019). Efficacy of dried *Phyllanthus amarus* leaf meal as an herbal feed additive on the growth performance, haematology and serum biochemistry of growing rabbits. *International Journal of Academic Research and Development*. 4(3): 97-104.
47. Alagbe, J.O. (2019). Growth response and bacteria count of broiler starter given *Delonix regia* leaf extract as a natural alternative to antibiotics. *Food and Nutrition: Current Research*. 2(3): 197 – 203.



This work is licensed under Creative Commons Attribution 4.0 License

To Submit Your Article Click Here:

Submit Manuscript

DOI:10.31579/2637-8914/258

Ready to submit your research? Choose Auctores and benefit from:

- fast, convenient online submission
- rigorous peer review by experienced research in your field
- rapid publication on acceptance
- authors retain copyrights
- unique DOI for all articles
- immediate, unrestricted online access

At Auctores, research is always in progress.

Learn more <https://auctoresonline.org/journals/nutrition-and-food-processing>