

Unusual Presentation of a Posterior Mediastinal Schwannoma Associated with Chest Pain, Cough, Hemoptysis and Dyspnea

Manouchehr Aghajanzadeh ^{1*}, Alreza Gafernejad², Hamidi Saidi Saidi.³, Ali Talebi⁴, Mahmood Khyabani ⁵, Omid Mosaffae Rad ¹, Zahra Sadin ⁶ and Mohaya Farzin ^{6,7}

¹Department of Thoracic Surgery, Guilan University of Medical Sciences, Rasht, Iran

²Department of Pulmonology, Guilan University of Medical Sciences, Rasht, Iran

³Department of Radiooncology, Guilan University of Medical Sciences, Rasht, Iran

⁴Department of pathology, Guilan University of Medical Sciences, Rasht, Iran

⁵Department of Radiology, Guilan University of Medical Sciences, Rasht, Iran

⁶Inflammatory Lung Diseases Research Center, Department of Internal Medicine, Razi Hospital, School of Medicine, Guilan University of Medical Sciences, Rasht, Iran.

⁷Department of physiology, Razi Clinical Research Development Center, Guilan University of Medical Sciences, Rasht, Iran

*Corresponding Author: Manouchehr Aghajanzadeh, Department of Thoracic Surgery, Guilan University of Medical Sciences, Rasht, Iran.

Received Date: August 22, 2024; Accepted Date: August 29, 2024; Published Date: September 06, 2024

Citation: Manouchehr Aghajanzadeh, Alreza Gafernejad, Hamidi Saidi Saidi, Ali Talebi, Mahmood Khyabani, et al, (2024), Unusual Presentation of a Posterior Mediastinal Schwannoma Associated with Chest Pain, Cough, Hemoptysis and Dyspnea, *J, Surgical Case Reports and Images*, 7(8); DOI:10.31579/2690-1897/213

Copyright: © 2024, Manouchehr Aghajanzadeh. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract

Neurogenic tumors (NT) are the most common of posterior mediastinal tumors and 80% of all posterior mediastinal tumors are (NT). They originate from the spinal cord, sympathetic ganglia, or peripheral nerve roots. A schwannoma is a slow-growing, encapsulated, and benign neurogenic tumor. Fewer than 9% of schwannomas are located in the mediastinum. Posterior mediastinal schwannomas originate from neural crest cells and typically from the intercostal nerves. Mediastinal schwannomas are often asymptomatic but may present with unusual symptoms as cough or dyspnea and hemoptysis because of pulmonary involvement. Although a definitive diagnosis is made by histopathology and immunohistochemical analysis and electron microscopy can be used to help its final diagnosis. We present a case of a benign posterior mediastinal schwannoma which presented in CXR and computed tomography (CT) imaging as a cystic lesion in a 58-year-old woman who admitted to our hospital with back pain, cough, hemoptysis and dyspnea.

Keywords: parenchyma preserving; hepatectomy; lesionectomy; metastatectomy; deep-seated liver lesions

Introduction

Several mediastinal classification systems have been discussed, which historically were based on no anatomic divisions on the lateral chest radiographs [1,2,3]. One of the classification systems includes perivascular (anterior), visceral (middle), and paravertebral (posterior) compartments [2,3]. Localization of a mediastinal lesion can help to better diagnosis and treatment. The paravertebral compartment is bound superiorly by the thoracic inlet, inferiorly by the diaphragm, anteriorly by the posterior boundary by pericardium, and posterior by lateral margin of the transverse processes and inner surface the chest wall [2,3]. Organs contains of posterior mediastinum include of descending aorta,

esophagus, thoracic duct, and vagus nerve. Neurogenic tumors of posterior mediastinum originate from the neural crest as peripheral, sympathetic, or paraganglionic elements, tumors of lymphatic, vascular, and mesenchymal origin may also occur [1,2,3]. Posterior mediastinal masses and lesion according to their origin and morphology can divide as neurogenic, esophageal, cystic lesion, extramedullary hematopoiesis, or lymphoma. 80% of posterior mediastinal masses are Neurogenic tumors [3,4,22]. There are two broad categories of neurogenic neoplasms: peripheral nerve sheath tumors such as schwannoma or neurofibroma which present as round or dumbbell lesions and the sympathetic ganglion

neoplasms such as ganglioneuroma, ganglioneuroblastoma and neuroblastoma which present as elongated masses involving 3 or more vertebral levels [2,3]. Other less common neoplasms in the differential include lymphoma, bone tumors, and metastases [2,3,4,5,6]. No neoplastic entities include spinal infections, cystic lesions (meningocele and neurenteric cyst), pancreatic pseudocyst and extramedullary hematopoiesis lesion, sympathetic ganglia. A schwannoma is a slow-growing, encapsulated, and benign neurogenic tumor. Fewer than 9% of schwannomas are located in the mediastinum. Posterior mediastinal schwannomas originate from neural crest cells and typically originate from the intercostal nerves [3,4,7,22]. Mediastinal schwannomas are often asymptomatic but may present with cough or dyspnea, hemoptysis and pain [2,3,4,5]. Although a definitive diagnosis is by histopathology, immunohistochemical analysis and electron microscopy can be used for definitive diagnosis [2,3,4,8,9,10]. Surgical resection is the primary treatment of choice in most neurogenic tumors, including the schwannoma [2,3,4,5,22]. A minimally invasive resection can use for resection [7,8,9,10,11]. The histology of mediastinal schwannoma is similar to that in other locations, presenting as an encapsulated tumors composed of spindle cells with typical variability in cellularity [11,12,13]. We present a case of a benign posterior mediastinal cystic schwannoma which present with cough, dyspnea, hemoptysis and chest pain and discovered by CXR and (CT-scan) in a 58-year-old woman and resect with right side posterolateral thoracotomy without any complications.

Case


A 58-year-old woman was referred to our hospital with a history of recurrent right side chest wall pain, cough, night sweating, hemoptysis

and dyspnea for 4-6 weeks. A physical examination revealed respiratory sounds were decreased on the middle right side of chest. His complete blood count, erythrocyte sedimentation rate, CRP blood urea nitrogen (BUN) level, and creatinine level were normal and liver function tests (AST, ALT, AKP and bilirubin) and anti-hydatid tests were normal. Poster anterior (PA) and lateral chest radiographs were performed. The PA chest radiograph (Figure 1) showed a well-defined 10- to 14-cm right paracardiac cystic mass located in the right middle zone of chest. The CT-scan showed a large well-defined hypo dense cystic mass measuring 112 mm is seen in superior segment of RUL. Sub segmental atelectasis also seen in right lung caused compressing. Left lung was normal. Figure 2(a,b,c,d,e,f,g,h,i).

In the CT-scan there was no invasion of the chest wall and others intrathoracic structures. CT images show no enlargement of lymph nodes and pleural effusion but lung paracardiac was involved. The testes for hydatid cyst were negative. Ultrasonography of abdomen was normal. A classic right posterolateral thoracotomy in six intercostal space was performed. Around of cystic lesion was walling off with wet sponge with normal Saline (Fig 3). In the aspiration of the lesion there was a hemorrhagic dense viscous fluid. The cystic mass was completely removed without complications and free margin. On the operation time, macroscopic examination show hemorrhagic dense viscous fluid and debris was observed in the center of the lesion (Fig 4) Microscopic examination showed an encapsulated mesenchymal neoplasm composed of short fascicles of bland looking spindle cell which are diffusely positive for S100 and GEAP on immunohistochemically diagnosis was a schwannoma tumor. The patient was discharged on the 6th postoperative day. The patient did not recurrence after 8 months of follow-up



Figure 1: CXR show right side lung cyst








Figure A,B Fig C,D Fig E,F,G

Figure 2: CT -scan A,B,C,D,E,F,G show solid cystic lesion



Figure 3: A,B,C,D show walling of around cystic lesion for prevention of spillage of fluid and debris of cystic masses




Figure 4: A,B,C pathology Microscopic examination showed an encapsulated mesenchymal neoplasm composed of short fascicles of bland looking spindle cell which are diffusely positive for S100 and GEAP on immunohistochemically diagnosis was a schwannoma tumor.



Figure 6: CXR 8 month postoperative

Discussion

The anterior border of the posterior mediastinum is pericardium and trachea, posterior border and lateral border is inner chest wall and the spine column (2,3) Organs contains of posterior mediastinum include of descending aorta, esophagus, thoracic duct, and vagus nerve (1,2,3,4). neurogenic tumors of posterior mediastinum originate from the neural crest as peripheral, sympathetic, or paraganglionic elements. tumors of lymphatic, vascular, and mesenchymal origin may also occur [1,2,3,4]. Posterior mediastinal masses and lesion according to their origin and morphology can divide as neurogenic, esophageal, cystic, extramedullary hematopoiesis, or lymphoma. 80% of posterior mediastinal masses are Neurogenic tumors 4. about 10- 20% of (2,3,4,7,8,22). Lesions are malignant [3,4]. neurofibroma, schwannoma, neurogenic sarcoma originate from peripheral nerve roots. ganglioneuroma, ganglioneuroblastoma, and neuroblastoma originated from aorticosympathetic paraganglia and paravertebral paraganglioma or from sympathetic ganglia in the Posterior mediastinum or rarely, meningocele or meningomyelocele originated from the intrathoracic spinal canal (1,2,3,4,5,9,10). Among mediastinal neurogenic tumors, schwannomas and neurofibromas arise from peripheral nerves and are more common in

adults, whereas ganglioneuromas and neuroblastomas arise from sympathetic ganglia and are more common in children [3,4,5,6,7,11,12]. The most common pediatric mediastinal neurogenic tumors are the ganglioneuroma and neuroblastoma (2,3,4,5). While the ganglioneuroma is a benign tumor, the neuroblastoma is highly malignant [2,3]. In the adult population, the neurofibroma and schwannoma are most common, with the schwannoma being the most common posterior mediastinal neurogenic neoplasm [2,3,4,6,8,9]. Schwannomas are often asymptomatic and incidentally detected during imaging. Schwannomas are mostly solid tumors, some of which may contain cystic degenerations or hemorrhages as our case. Rarely they present as a completely cystic lesion [8,9,10,11]. Schwannomas originating from the nerve roots are located in the paravertebral area, and in the anterior vertebral corpus and neural foramen. PA and lateral chest x-rays are usually the first test performed, as our case [2,3,4,5,15,16,22]. but a CT scan with intravenous contrast is the most valuable test for mediastinal lesions. In this case after CXR we used CT-scan [2,3,4,5,8], Posterior mediastinum masses usually compress the lung and produce Symptoms may occur due to the compression of adjacent mediastinal structures, such as the airway, esophagus, heart, and great vessels [2,3,5,15,16]. Although they are rarely

malignant, they may cause respiratory symptoms, such as stridor, dyspnea, hemoptysis, and cough, or gastrointestinal symptoms, such as dysphagia, in the case of invasion to local structures [3,4,5,18]. Our case present with cough, dyspnea, chest pain and hemoptysis. Symptoms of posterior mediastinum neurogenic tumors include compression of the airway, esophagus, heart, and great veins. Malignant tumors of this area can of invasion into the tracheobronchial tree, spinal canal, lungs, esophagus, superior vena cava, pleura, and chest wall, these invasion can produce cough, stridor, dyspnea, hemoptysis, dysphagia, pleural effusion, and superior vena cava syndrome and hoarseness, neuropathic pain, diaphragmatic paralysis, or Horner's syndrome and Paraneoplastic syndromes. In our case presentation was due to compressions to adjacent organs involvement [2,3,4,5,9,18,19]. The initial workup of a posterior mediastinal mass involves radiographic evaluation, with the majority of asymptomatic masses discovered incidentally on posteroanterior (PA) and lateral radiographs [2,3,4,14]. Additional radiologic findings include enlargement of the neural foramina (creating a dumbbell shaped lesion), scalloping of posterior vertebral bodies, erosion of the ribs, pleural effusion, and scoliosis. Disruption of the azygoesophageal recess is a nonspecific finding seen in both middle and posterior mediastinal masses [3,16,17]. Posterior masses generally have sharp margins due to their interface with the lung, CT imaging of the mass is useful in determining the exact location and its relationship to adjacent structures and may aid in differentiating tissue densities between cystic, vascular, and solid masses [2,3,4,6,7]. Due to high contrast resolution and multiplanar capability, MRI is the preferred modality of imaging as it provides better evaluation regarding the nature and extent of intraspinal involvement towards the neural and vertebral foramen [3,17]. The limitations of MRI compared to CT are the limited radiographic evaluation of calcifications and poorer spatial resolution [2,3,4,16,17] therefore, while an MRI is preferable it is not always necessary, we do not MRI in our case [2,3,21]. Recent studies have offered ultrasound guided FNA as a viable diagnostic modality for accessible mediastinal lesions [2,3,23] allowing for formal diagnosis without the potential complications from more invasive procedures such as CT-guided biopsy, mediastinoscopy, or video-assisted thoracoscopic surgery which offer no benefit in the management of these tumors [2,3]. Because of suspicion of hydatid cyst our radiologist team do not use FNA [23,24], hydatid cyst in IRAN are endemic [22]. PET scans have shown the ability to distinguish between malignant peripheral nerve sheath tumors and neurofibromas with high accuracy but are not as helpful in distinguishing between benign and malignant peripheral nerve sheath tumors [2,3,16,17]. Primary treatment of schwannoma and most neurogenic tumors are Surgical resection [2,3,4,5,19]. schwannomas may originate from the vagus, phrenic, or any part of the intercostal nerve and removed of tumor should be removed by a saving the involved nerve [2,3,18,19], Radiation therapy may be used postoperatively to control residual disease in malignant schwannomas, but its benefit is unknown [2,3,4,5]. No known chemotherapeutic regimens are effective against these tumors [3,5,6,7]. Since serologic markers are absent and characteristic imaging abnormalities are variable, tissue pathology and immunohistochemistry are required for a diagnosis [11,12,13]. Schwannomas are composed of spindle cells with twisted nuclei, amphophilic cytoplasm, and rare mitoses [2,3,11,12,13]. Patients with neurofibromatosis are likely to display a variant form called plexiform schwannoma [12,13]. Another variant, the melanocytic schwannoma, has a pronounced brownish cytoplasmic pigment and malignant potential [11,12,13]. The malignant schwannoma, the most dangerous variant, is a soft gray-pinkish tumor with central necrosis and microscopically

consists of sheets of pleomorphic spindle cells with numerous mitotic figures and necrotic areas [3,11,12,13]. Malignant schwannomas may also exhibit a variety of cellular components such as clusters of epithelial cells; mucin-secreting glands; and even mesenchymal features such as bone, cartilage, or skeletal muscle [3,10,11,12,13]. Patients who undergo resection of a benign schwannoma are generally followed up to monitor wound healing and resumption of daily activities. Routine follow-up of patients who undergo curative resection of a malignant neoplasm has not demonstrated a survival benefit in randomized controlled trials due to the variety and infrequency of malignant mediastinal tumors [2,3,12,15,16]. Overall, patients with benign schwannoma have excellent survival following complete resection, whereas those with malignant tumors have a poorer prognosis [2,3,5,22,23,24].

Acknowledgment

The authors wish to thank Roya Moghadam, Chraghalizadeh Executive Assistant in the Department of Thoracic Nursing at the Arya private hospital. for her and his skillful expertise in preparation of the figures for publication

We acknowledge the great work of our medical officer Dr Ahmad Reshad Wajed, and our histotechnologists; Mohammad Ibrahim Wali Zada, Esmatullah Ramzi, and Adil Muhammedi.

References

1. Carter BW, Tomiyama N, Bhora FY, et al. (2014). A modern definition of mediastinal compartments. *J Thorac Oncol* 2014;9: S97-S101
2. Carter BW, Benveniste MF, Madan R, et al. (2017). ITMIG Classification of Mediastinal Compartments and Multidisciplinary Approach to Mediastinal Masses. *Radiographics*; 37:413-436
3. John Matthew Archer, Jitesh Ahuja, Chad D. Strange, Girish S. Shroff, Gregory W. Gladish, Bradley S. Sabloff, Mylene T. Truong Multimodality imaging of mediastinal masses and mimics, *MEDIASTINUM*, Vol 7 (September 25, 2023)
4. Mubashir M, Salam A, Sonawalla A, Fatima H, Fatimi SH. (2017). Rare presentation of a posterior mediastinal cystic schwannoma as a large pleural effusion. *Cureus*. 9:e1558.
5. Ruchi Amin, Brett H. (2015). Waibel, An Unusual Presentation of a Posterior Mediastinal Schwannoma Associated with Traumatic Hemothorax, *Case Reports in Surgery* First published.
6. Septiardi Husodo, Farah Fatma Wati. (2023). Mediastinal schwannoma: A case report, *International Journal of Surgery Case Reports*, 108564
7. Zhu D, Hong P, Chen S, Fan Y, Zhang Z. et al. (2018). Lower posterior mediastinal benign schwannoma successfully resected with retroperitoneoscopy using a transdiaphragmatic approach: A case report. *Thorac Cancer*. 9:1087-1089.
8. Mubashir M, Salam A, Sonawalla A, Fatima H, Fatimi SH. (2017). Rare presentation of a posterior mediastinal cystic schwannoma as a large pleural effusion. *Cureus*;9:e1558.
9. Amin R, Waibel BH. (2015). An unusual presentation of a posterior mediastinal schwannoma associated with traumatic hemothorax. *Case Rep Surg*.175645.

10. Alkan S, Dadaş B. (2007). Boyunda dev schwannoma olgusu. *Şişli Etfal Hastanesi Tıp Bülteni*. 1:47-49.
11. Aslı Tanrıvermiş Sayıt1. (2020). Muzaffer Elmalı1 Rare presentation of a posterior mediastinal schwannoma *Cukurova Medical Journal*;45(2):760-762
12. Sugita S, Aoyama T, Emori M *et al.* (2021). Assessment of H3K27me3 immunohistochemistry and combination of NF1 and p16 deletions by fluorescence in situ hybridization in the differential diagnosis of malignant peripheral nerve sheath tumor and its histological mimics. *Diagn. Pathol.*16; 79.
13. Asano N, Yoshida A, Ichikawa H *et al.* (2017). Immunohistochemistry for trimethylated h3k27 in the diagnosis of malignant peripheral nerve sheath tumours. *Histopathology* ; 70; 385–393.
14. Mito JK, Qian X, Doyle LA, Hornick JL, Jo VY. (2017). Role of histone h3k27 trimethylation loss as a marker for malignant peripheral nerve sheath tumor in fine-needle aspiration and small biopsy specimens. *Am. J. Clin. Pathol.* 148; 179–189.
15. Mohammad Reza Asgary, Manouchehr Aghajanzadeh, Hosein Hemmati, Morteza Jafari. (2016). Epidemiology Of Mediastinal Tumors During Six Years (2006-2012) In Rasht City. *Acta Medica Mediterranea*, 32: 1243
16. Strollo DC, Rosado-de-Christenson ML, Jett JR. (1997). Primary mediastinal tumors: part II. Tumors of the middle and posterior mediastinum. *Chest*; 112:1344-1357
17. Bredella MA, Torriani M, Hornicek F, et al. (2007). Value of PET in the assessment of patients with neurofibromatosis type 1. *AJR Am J Roentgenol*; 189:928-35
18. Broski SM, Johnson GB, Howe BM, et al. (2016). Evaluation of (18)F-FDG PET and MRI in differentiating benign and malignant peripheral nerve sheath tumors. *Skeletal Radiol* ;45:1097-1105.
19. Wilson LD, Detterbeck FC, Yahalom J. (2007). Clinical practice. Superior vena cava syndrome with malignant causes. *N Engl J Med*; 356:1862-1869.
20. Okamatsu C, London WB, Naranjo A *et al.* (2009). Clinicopathological characteristics of ganglioneuroma and ganglioneuroblastoma: a report from the ccg and cog. *Pediatr. Blood Cancer*; 53; 563–569.
21. Ocazonez D, Shroff GS, Vargas D, et al. (2017). Imaging of Intrathoracic Paragangliomas. *Semin Ultrasound CT MR*;38:584-593.
22. Raptis CA, McWilliams SR, Ratkowski KL, et al. (2018). Mediastinal and Pleural MR Imaging: Practical Approach for Daily Practice. *Radiographics*; 38:37-55.
23. Manouchehr Aghajanzadeh, Sna Khajeh Jahromi. (2014). Rasool Hassanzadeh, Hannan Ebrahimi. Posterior Mediastinal Cyst. January, *Archives of Iranian Medicine*; 17(1):95-96.
24. Benz M. R., Czernin J., Dry S. M. et al. (2010). Quantitative F18-fluorodeoxyglucose positron emission tomography accurately characterizes peripheral nerve sheath tumors as malignant or benign, *Cancer*. 116, 451–458.
25. Pakseresht K., Reddymasu S. C., Oropeza-Vail M. M., Fan F., and Olyae M. (2011). Mediastinal schwannoma diagnosed by endoscopic ultrasonography-guided fine needle aspiration cytology, *Case Reports in Gastroenterology*. (2011) 5, no. 2, 411–415



This work is licensed under Creative Commons Attribution 4.0 License

To Submit Your Article Click Here:

Submit Manuscript

DOI:10.31579/2690-1897/213

Ready to submit your research? Choose Auctores and benefit from:

- fast, convenient online submission
- rigorous peer review by experienced research in your field
- rapid publication on acceptance
- authors retain copyrights
- unique DOI for all articles
- immediate, unrestricted online access

At Auctores, research is always in progress.

Learn more <https://auctoresonline.org/journals/journal-of-surgical-case-reports-and-images>