

The Relationship between Physical Activity, Women's Overall Health, and well-being

Onur Oral ^{1*}, Pramila Thapa ², Pinar Tatlıbal ³, Mumtaz Enser ⁴, Evangelia Stavropoulou ⁵

¹Ege University, Faculty of Sports Sciences, Izmir, Turkey.

²Life Skill Education Institutes/Yeti Health Science Academy, Katmandu, Nepal.

³Dokuz Eylul University, Faculty of Necat Hepkon Sport Sciences, Izmir, Turkey.

⁴Dokuz Eylul University, Institute of Social Sciences, Department of Philosophy, Izmir, Turkey

⁵Master of Research in Information and Communication, University of Lorraine, France

***Corresponding Author:** Onur Oral, University of Ege, Faculty of Sports Sciences, Department of Health Sciences and Sports, Izmir, Turkey.

Received Date: 21 October 2024 | **Accepted Date:** 04 November 2024 | **Published DATE:** 10 December 2024

Citation: Onur Oral, Pramila Thapa, Pinar Tatlıbal, Mumtaz Enser, Evangelia Stavropoulou, (2024), The Relationship between Physical Activity, Women's Overall Health, and well-being, *J. Endocrinology and Disorders*, 8(6): DOI:10.31579/2640-1045/198

Copyright: © 2024, Onur Oral. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Abstract

Background:

This review article suggests that there are many positive effects of physical activity and exercise on women's overall health. It is widely acknowledged that regular exercise can have a beneficial impact on a number of areas, including cardiovascular health, weight management, bone health, mental well-being, cancer risk and longevity.

Materials and Methods:

In order to gain a comprehensive understanding of the topic, a literature search was conducted using various databases and keywords related to physical activity, longevity, women's health, quality of life, healthy aging, and obesity. The search was limited to peer-reviewed articles published in English between 1994 and 2024. Animal model studies were excluded to focus on human research. The criteria for inclusion in this review were that the studies examined the association between weight control management and overall healthy aging, were published in a peer-reviewed journal, and were written in English. After careful consideration, a total of 38 studies were selected for inclusion.

Results:

Regular exercise is of great importance for the prevention and treatment of overweight and obesity, as well as for the management of numerous chronic diseases that have become prevalent in recent times. Furthermore, it plays a pivotal role in maintaining regular metabolic processes. Integrating regular physical activity into one's daily life, in conjunction with a natural and balanced diet, can facilitate the maintenance of a healthy lifestyle and the maintenance and improvement of overall health. It would therefore be beneficial for women of all ages to consider integrating physical activity into their daily routines to facilitate the enjoyment of these advantages and contribute to the development of a healthier lifestyle.

Conclusion:

The scientific evidence is compelling in supporting the promotion of regular physical activity as a fundamental aspect of women's health. It is imperative that healthcare providers and policymakers prioritize initiatives that encourage and facilitate physical activity among women of all ages. By doing so, we can move toward a future where a greater proportion of women can enjoy the numerous health benefits of an active lifestyle, which will lead to improved quality of life and increased longevity.

Key words: physical activity; longevity; women's health; quality of life; healthy aging; obesity

Introduction

Physical activity and exercise are integral components of maintaining overall health and well-being for women. Engaging in regular physical

activity not only aids in weight management but also significantly enhances cardiovascular health, mitigates the risk of chronic diseases,

improves mental well-being, and promotes longevity Brown, Burton & Rowan, 2007; Eyler et al., 1998). This systematic literature review aims to investigate the multifaceted effects of physical activity and exercise on women's health by examining a variety of research studies in this domain.

It is well documented that regular physical activity has a beneficial effect on cardiovascular health in women. It is widely accepted that aerobic exercise, including resistance training, especially brisk walking, running, and cycling, has the potential to strengthen the heart muscle, improve blood circulation, reduce the risk of heart disease, and be an effective method of weight control in the prevention and treatment of obesity (Hu et al., 2001; Myers et al., 2002; Oral et al., 2023). Research consistently indicates that women who engage in regular physical activity exhibit a significantly reduced risk of developing hypertension, stroke, and heart attacks compared to their sedentary counterparts (Haapanen et al., 1997; Oguma & Shinoda-Tagawa, 2004). Furthermore, exercise positively influences lipid profiles by increasing high-density lipoprotein (HDL) cholesterol, often referred to as "good" cholesterol, while simultaneously reducing low-density lipoprotein (LDL) cholesterol, known as "bad" cholesterol (Chandra et al., 2014).

It is widely acknowledged that physical activity and exercise play an important role in maintaining a healthy weight in women. In addition to burning calories, regular exercise has been shown to positively affect metabolic processes and have beneficial effects on the cardiometabolic and immune systems. In light of these findings, it seems reasonable to suggest that physical activity may contribute to more effective weight management and metabolic improvements, potentially providing critical support for longevity, quality of life, and healthy aging processes in women. (Donnelly et al., 2009; Williams et al., 2015; Rezaee et al., 2022). It would be advantageous for women to implement a balanced exercise regimen that incorporates both aerobic activities, such as walking and cycling, and strength training, specifically anaerobic exercises, in order to achieve their desired weight and prevent weight gain over time (Oral et al., 2024). Furthermore, physical activity contributes to improved body composition by reducing fat mass and increasing lean muscle mass, which positively impacts overall physical appearance and self-esteem (Aldhahi et al., 2022; Sattar, Khan & Iqbal, 2020).

In addition to physical health benefits, regular exercise is associated with improved mental well-being among women. Studies indicate that physical activity can alleviate symptoms of anxiety and depression, promoting a more positive mood and overall psychological resilience (Chu et al., 2009; Mandolesi, et al., 2018; Singh et al., 2023; Rezaee, et al., 2022). The release of endorphins during exercise contributes to feelings of happiness and relaxation, which can be especially beneficial for women managing stress or facing life challenges (Klonoff, Annechild & Landrine, 1994; Sran et al., 2021).

Materials and Methods:

A comprehensive search of the literature was undertaken using a range of databases, including the US National Library of Medicine (PubMed), Scopus, EBSCO, MEDLINE, DRJI, (Directory of Research Journal Indexing), Embase, Web of Science, Google Scholar, and SportDiscus. In order to gain a comprehensive understanding of the subject matter, a number of keywords were selected, including "physical activity", "longevity", "women's health", "quality of life", "healthy aging", and "obesity". In addition, relevant literature was also sourced from searching for articles in reference lists derived from the data searches. The search was limited to peer-reviewed articles published in English between 1994 and 2024, to focus the results. To gain a comprehensive understanding of the relationship between metabolic health, obesity, longevity, and various systemic conditions, studies that involved animal models were excluded from the research. To be included in this review, studies had to meet some criteria. Firstly, they had to examine the relationship between physical activity and women's overall health and well-being. Secondly, they had to have been published in a peer-reviewed

journal. Thirdly, they had to be available in English. After careful consideration, a total of 38 studies were selected for inclusion in this review.

Results and Discussion:

The role of exercise in weight management is well-documented. Donnelly et al. (2013) and Fogelholm and Kukkonen-Harjula (2000) both underscore the importance of regular physical activity in preventing weight gain and aiding in weight loss. The overall impression that can be gathered from these studies is that a training program which incorporates not only aerobic but also a combination of aerobic and resistance training might be an effective approach for women seeking to maintain a healthy weight. Such a program could potentially contribute to reducing the risk of developing obesity-related conditions such as obesity, type 2 diabetes, cardiovascular disease and hypercholesterolemia (Oral, Tatlibal & Stavropoulou, 2021).

Exercise is crucial for cardiovascular health, as it helps improve heart function, reduce blood pressure, and lower cholesterol levels. Kokkinos and Myers (2010) provide evidence that regular physical activity can lead to significant improvements in cardiovascular outcomes, reducing the risk of heart disease. Warburton et al. (2006) also highlight the broad spectrum of cardiovascular benefits associated with regular exercise, emphasizing its role in enhancing overall cardiovascular health.

One of the critical areas in which exercise plays a significant role is in the prevention and management of osteoporosis. In a 2009 study, Schmitt, Schmitt and Dören emphasized the efficacy of weight-bearing and resistance exercises in increasing bone mineral density (BMD), thereby reducing the risk of fractures. Similarly, Singh (2015) and Manaye et al. (2023) underscored the importance of regular physical activity, particularly high-impact and resistance training, for maintaining bone health in women, particularly postmenopausal women who are at an elevated risk for osteoporosis.

Physical activity has also been linked to a reduced risk of certain cancers. McTiernan et al. (2019). conducted a research study, concluding that regular exercise is associated with a lower risk of breast cancer. Lynch, Neilson and Friedenreich, (2011) further support this finding, indicating that physical activity interventions can play a preventive role in various types of cancer, likely due to its effects on hormone regulation and immune function.

The social aspects of physical activity are also important for women's health. Hawkey, Thisted and Cacioppo (2009) and Tsuji et al. (2020) discuss how engaging in group exercises or sports can enhance social interactions, reduce feelings of isolation, and improve overall quality of life. These social benefits are particularly relevant for women, who may experience unique social and psychological challenges.

Exercise has been shown to have substantial benefits for mental health, including the management of clinical depression. Craft and Perna (2004) and Rebar et al. (2015). provide evidence that regular physical activity interventions can significantly reduce symptoms of depression, offering a non-pharmacological treatment option that can be particularly beneficial for women. The psychological benefits of exercise also extend to improved self-esteem and overall mental well-being, as demonstrated by Haugen, Ommundsen, and Seiler (2013). Their findings indicated a positive correlation between physical activity interventions and self-esteem in women.

In light of the substantial contributions that physical activity makes to overall wellness, it would be advantageous to have a comprehensive framework in place to support the implementation of effective physical activity programs for women. This framework should take into account the specific approaches that are required to account for individual preferences, as well as the barriers and facilitators that may influence the success of such programs. When designing effective exercise programs

for women, it would be beneficial to consider this framework in order to maximize adherence and benefits (Larsen et al., 2015; Rebar et al., 2016).

Conclusion:

There is increasing evidence that regular physical activity, especially aerobic exercise, can have a positive effect on cardiovascular health in women. Scientific evidence has been shown in research studies that activities such as brisk walking, running, or cycling can help strengthen the heart muscle, improve blood flow, and reduce the risk of heart disease. The fact that there is a correlation between being physically active and having a lower risk of developing high blood pressure, stroke, or heart attack is an important gain for women's overall health and healthy aging. It is also thought that exercise can help improve cholesterol levels by increasing the amount of good cholesterol (HDL) and decreasing the amount of bad cholesterol (LDL). It is also stated that physical activity can play an important role in weight management and is an effective method for preventing obesity. It is thought that regular exercise can help burn calories, build muscle, and speed up metabolism, potentially helping with weight loss and maintenance. It may be beneficial for women to consider combining aerobic exercise with strength training to reach their desired weight and prevent weight gain over time. It is thought that women may be more prone to developing osteoporosis and in this respect, it is possible that participating in weight-bearing exercises may help to improve bone density and reduce the risk of this condition. Considering starting these exercises at a younger age may be beneficial for the health of the skeletal system in later life.

There seems to be increasing evidence that regular physical activity interventions can have a positive effect on mental health, including reducing symptoms of depression, anxiety, and stress. Group exercise activities may also have the potential to promote social interaction and improve mental well-being, as exercise can help stimulate the release of endorphins, which can contribute to a better mood. It is also suggested that exercise may help improve body composition by reducing fat mass and increasing lean muscle mass, which may lead to improvements in physical appearance and self-esteem.

There is some significant evidence from clinical research studies to suggest that regular exercise, particularly at moderate to vigorous intensity, may have the potential to reduce the risk of certain types of cancer in women, including breast, colon, and endometrial cancer. Although the exact mechanisms behind this protective effect are not fully understood, it is believed that exercise helps regulate hormone levels, reduce inflammation, and improve immune function, all of which contribute to cancer prevention.

It has also been suggested that physical activity may contribute to a longer life expectancy in women; physically active women may have a lower risk of premature death compared to those who are sedentary. In addition to these benefits, physical activity may also help with weight management and cardiovascular health, reduce the risk of chronic disease, and increase longevity. It is thought that regular exercise may help reduce the risk of chronic diseases, such as heart disease, stroke, diabetes, and some cancers, which are leading causes of death, and it may be associated with promoting healthy aging because exercise helps maintain muscle strength, flexibility, and cognitive function. All of these positive health benefits of regular physical activity interventions may help women maintain an active and independent lifestyle as they age, increasing their chances of longevity.

In conclusion, there is a great deal of evidence to suggest that physical activity can have a positive impact on women's health. It would be remiss of us not to mention the numerous benefits of regular exercise. These include improvements to cardiovascular health and weight management, as well as enhancements to bone density and mental well-being. It would be beneficial for women to consider integrating various forms of physical activity into their daily routines, to maximize health outcomes and promote a higher quality of life.

Acknowledgment:

We would like to express our special thanks to Dr. George N. NOMIKOS for his very successful contribution to the literature research process and unique academic support in the publication during the process of this review article.

Conflict of interest:

The author certifies that there is no conflict of interest with any financial organization regarding the material discussed in the manuscript.

Funding:

The author certifies that there is no funding from any financial organization regarding the material discussed in the manuscript or contributions:

Author contributions:

All the authors read and approved the final version of the manuscript.

References:

1. Aldhahi, M. I., Al Khalil, W. K., Almutiri, R. B., Alyousefi, M. M., Alharkan, B. S., & AnNasban, H. (2022). Effect of weight self-stigma and self-esteem on aerobic exercise capacity in adult women with different body compositions. *International Journal of Environmental Research and Public Health*, 19(2):873.
2. Brown, W. J., Burton, N. W., & Rowan, P. J. (2007). Updating the evidence on physical activity and health in women. *American journal of preventive medicine*, 33(5):404-411.
3. Chandra, K. S., Bansal, M., Nair, T., Iyengar, S. S., Gupta, R., Manchanda, S. C., ... & Gulati, S. (2014). Consensus statement on management of dyslipidemia in Indian subjects. *Indian heart journal*, 66:1.
4. Chu, I. H., Buckworth, J., Kirby, T. E., & Emery, C. F. (2009). Effect of exercise intensity on depressive symptoms in women. *Mental Health and Physical Activity*, 2(1), 37-43.
5. Craft, L. L., & Perna, F. M. (2004). The benefits of exercise for the clinically depressed. *Primary care companion to the Journal of clinical psychiatry*, 6(3):104.
6. Donnelly, J. E., Blair, S. N., Jakicic, J. M., Manore, M. M., Rankin, J. W., & Smith, B. K. (2009). Appropriate physical activity intervention strategies for weight loss and prevention of weight regain for adults. *Medicine & Science in Sports & Exercise*, 41(2): 459-471.
7. Donnelly, J. E., Honas, J. J., Smith, B. K., Mayo, M. S., Gibson, C. A., Sullivan, D. K., ... & Washburn, R. A. (2013). Aerobic exercise alone results in clinically significant weight loss for men and women: midwest exercise trial 2. *Obesity*, 21(3):219-228.
8. Eyler, A. A., Brownson, R. C., King, A. C., Brown, D., Donatelle, R. J., & Heath, G. (1998). Physical activity and women in the United States: An overview of health benefits, prevalence, and intervention opportunities. *Women & health*, 26(3):27-49.
9. Haapanen, N., Miilunpalo, S., Vuori, I., Oja, P., & Pasanen, M. (1997). Association of leisure time physical activity with the risk of coronary heart disease, hypertension and diabetes in middle-aged men and women. *International journal of epidemiology*, 26(4): 739-747.
10. Fogelholm, M., & Kukkonen-Harjula, K. (2000). Does physical activity prevent weight gain—a systematic review. *Obesity reviews*, 1(2):95-111.
11. Haugen, T., Ommundsen, Y., & Seiler, S. (2013). The relationship between physical activity and physical self-esteem in adolescents: the role of physical fitness indices. *Pediatric exercise science*, 25(1):138-153.

12. Hawkey, L. C., Thisted, R. A., & Cacioppo, J. T. (2009). Loneliness predicts reduced physical activity: cross-sectional & longitudinal analyses. *Health psychology*, 28(3):354.
13. Hu, F. B., Stampfer, M. J., Solomon, C., Liu, S., Colditz, G. A., Speizer, F. E., ... & Manson, J. E. (2001). Physical activity and risk for cardiovascular events in diabetic women. *Annals of internal medicine*, 134(2):96-105.
14. Klonoff, E. A., Annechild, A., & Landrine, H. (1994). Predicting exercise adherence in women: the role of psychological and physiological factors. *Preventive Medicine*, 23(2):257-262.
15. Kokkinos, P., & Myers, J. (2010). Exercise and physical activity: clinical outcomes and applications. *Circulation*, 122(16):1637-1648.
16. Larsen, B. A., Noble, M. L., Murray, K. E., & Marcus, B. H. (2015). Physical activity in Latino men and women: facilitators, barriers, and interventions. *American Journal of Lifestyle Medicine*, 9(1):4-30.
17. Lynch, B. M., Neilson, H. K., & Friedenreich, C. M. (2011). Physical activity and breast cancer prevention. *Physical activity and cancer*, 13-42.
18. Manaye, S., Cheran, K., Murthy, C., Bornemann, E. A., Kamma, H. K., Alabbas, M., ... & Franchini, A. P. A. (2023). The role of high-intensity and high-impact exercises in improving bone health in postmenopausal women: a systematic review. *Cureus*, 15(2).
19. Mandolesi, L., Polverino, A., Montuori, S., Foti, F., Ferraioli, G., Sorrentino, P., & Sorrentino, G. (2018). Effects of physical exercise on cognitive functioning and wellbeing: biological and psychological benefits. *Frontiers in psychology*, 9:509.
20. McTiernan, A. N. N. E., Friedenreich, C. M., Katzmarzyk, P. T., Powell, K. E., Macko, R., Buchner, D., ... & Piercy, K. L. (2019). Physical activity in cancer prevention and survival: a systematic review. *Medicine and science in sports and exercise*, 51(6):1252.
21. Myers, J., Prakash, M., Froelicher, V., Do, D., Partington, S., & Atwood, J. E. (2002). Exercise capacity and mortality among men referred for exercise testing. *New England journal of medicine*, 346(11):793-801.
22. Oguma, Y., & Shinoda-Tagawa, T. (2004). Physical activity decreases cardiovascular disease risk in women: review and meta-analysis. *American journal of preventive medicine*, 26(5):407-418.
23. Oral, O., Rezaee, Z., Nomikos, G., Nomikos, N., & Rashidlamir, A. (2023). The impact of both resistance and aerobic training on weight control management. *Scientific Chronicles/Epistimonika Chronika*, 28(2).
24. Oral, O., Rezaee, Z., Enser, M., Nomikos, N.G. (2024). The Effect of Regular Exercise on Women's Overall Health. *Ortho Res Online J.* 10(5). OPROJ. 000748.
25. Oral, O., Tatlibal, P., & Stavropoulou, E. (2021). A Narrative Review of the Impact of Aerobic Training on the Prevention and Treatment of Obesity-Related Hypertension. *Ortho Res Online J*, 8(3).
26. Rebar, A. L., Stanton, R., Geard, D., Short, C., Duncan, M. J., & Vandelanotte, C. (2015). A meta-meta-analysis of the effect of physical activity on depression and anxiety in non-clinical adult populations. *Health psychology review*, 9(3):366-378.
27. Rebar, A. L., Dimmock, J. A., Jackson, B., Rhodes, R. E., Kates, A., Starling, J., & Vandelanotte, C. (2016). A systematic review of the effects of non-conscious regulatory processes in physical activity. *Health psychology review*, 10(4):395-407.
28. Ross, R., Blair, S. N., Arena, R., Church, T. S., Després, J. P., Franklin, B. A., & Wisløff, U. (2016). Importance of assessing cardiorespiratory fitness in clinical practice: a case for fitness as a clinical vital sign: a scientific statement from the American Heart Association. *Circulation*, 134(24):653-699.
29. Sattar, S., Khan, S., & Iqbal, S. (2020). Impact of self-esteem and body image on sports participation of female athletes. *THE SKY-International Journal of Physical Education and Sports Sciences (IJPESS)*, (1):65-80.
30. Schmitt, N. M., Schmitt, J., & Dören, M. (2009). The role of physical activity in the prevention of osteoporosis in postmenopausal women—an update. *Maturitas*, 63(1), 34-38.
31. Singh, B., Olds, T., Curtis, R., Dumuid, D., Virgara, R., Watson, A., ... & Maher, C. (2023). Effectiveness of physical activity interventions for improving depression, anxiety and distress: an overview of systematic reviews. *British journal of sports medicine*, 57(18), 1203-1209.
32. Singh, M. A. F. (2015). Exercise and bone health. *Nutrition and bone health*, 505-542.
33. Sran, S. K., Vats, P., & Wadhawan, P. (2021). Effect of exercise on life satisfaction and happiness. *Indian Journal of Health and Wellbeing*, 12(1), 79-82.
34. Thune, I., Brenn, T., Lund, E., & Gaard, M. (1997). Physical activity and the risk of breast cancer. *New England Journal of Medicine*, 336(18), 1269-1275.
35. Tsuji, T., Kanamori, S., Saito, M., Watanabe, R., Miyaguni, Y., & Kondo, K. (2020). Specific types of sports and exercise group participation and socio-psychological health in older people. *Journal of sports sciences*, 38(4), 422-429.
36. Warburton, D. E., Nicol, C. W., & Bredin, S. S. (2006). Health benefits of physical activity: the evidence. *Cmaj*, 174(6), 801-809.
37. Williams, R. L., Wood, L. G., Collins, C. E., & Callister, R. (2015). Effectiveness of weight loss interventions—is there a difference between men and women: a systematic review. *obesity reviews*, 16(2), 171-186.
38. Z. Rezaee, S. Javaheri, A. Rashidlamir, O. Oral, E. Stavropoulos. (2022) Effects of BFR training in serum NRG-1 and IL-6 in overweight postmenopausal women. *Scientific Chronicles* 2022; 27(4): 541-554.



This work is licensed under Creative Commons Attribution 4.0 License

To Submit Your Article Click Here:

Submit Manuscript

DOI:10.31579/2640-1045/198

Ready to submit your research? Choose Auctores and benefit from:

- fast, convenient online submission
- rigorous peer review by experienced research in your field
- rapid publication on acceptance
- authors retain copyrights
- unique DOI for all articles
- immediate, unrestricted online access

At Auctores, research is always in progress.

Learn more <https://auctoresonline.org/journals/endocrinology-and-disorders>