

Autoimmune Hemolytic Anemia Secondary to *Mycoplasma pneumoniae* Infection in a Child with Chromosomal Abnormality: Case Report

Ayman Gouda*, Mahmoud Kamal Moustafa Ahmed, AnasAlwogud A. Abdelmogheth, Alddai Alniar, Ayisha Hussain, Shatha Alawneh, Hanan Abouelkheir

Al Qassimi Women's and Children's Hospital, Sharjah, United Arab Emirates, PICU department

*Corresponding Author: Ayman Gouda, Al Qassimi Women's and Children's Hospital, Sharjah, United Arab Emirates, PICU department

Received Date: November 24, 2025 | Accepted Date: December 15, 2025 | Published Date: December 18, 2025

Citation: Ayman Gouda, Mahmoud K M Ahmed, AnasAlwogud A. Abdelmogheth, Alddai Alniar, et al (2025), Autoimmune Hemolytic Anemia Secondary to *Mycoplasma pneumoniae* Infection in a Child with Chromosomal Abnormality: Case Report, *International Journal of Clinical Case Reports and Reviews*, 31(5); DOI:10.31579/2690-4861/1001

Copyright: © 2025, Ayman Gouda. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Abstract:

Mycoplasma pneumoniae is a common cause of respiratory infections in children and is well recognized for its potential extrapulmonary complications, including autoimmune hemolytic anemia (AIHA) [1,2]. Cold agglutinin disease represents a rare but severe form of AIHA associated with *Mycoplasma* infection and may lead to life-threatening hemolysis [3,7]. We report the case of a 7-year-old female with a pathogenic mosaic gain of chromosome 12p13 and multiple medical comorbidities who developed rapidly progressive cold agglutinin-mediated AIHA secondary to *Mycoplasma pneumoniae* infection. Her clinical course was marked by sudden hemodynamic collapse, profound anemia with a hemoglobin nadir of 2.7 g/dL, and an unusual presentation with bradycardia [4,5]. Diagnostic evaluation confirmed immune-mediated hemolysis with a positive direct antiglobulin test for complement, positive cold agglutinin testing, and positive *Mycoplasma* IgM serology [6]. Management included aggressive supportive care, packed red blood cell transfusions, pulse corticosteroid therapy, and targeted antibiotic treatment with azithromycin [2,7]. The patient showed progressive clinical and hematological recovery.

Key words: mycoplasma pneumoniae; autoimmune hemolytic anemia; cold agglutinin disease; chromosomal Abnormality; Pediatrics

Introduction

Mycoplasma pneumoniae is a frequent cause of community-acquired pneumonia in children and adolescents, accounting for up to 40% of pediatric cases [1,2]. Extrapulmonary manifestations may occur in 25–50% of hospitalized patients and include neurological, dermatological, cardiovascular, and hematological complications such as autoimmune hemolytic anemia (AIHA) [1,3]. Elevated cold agglutinin titers are detected in up to 50–60% of infected patients; however, clinically significant hemolysis remains rare [6,12]. Severe cold agglutinin-mediated AIHA can lead to rapid hemodynamic deterioration and requires early recognition and aggressive management [7,8].

Case Presentation

A 7-year-old female with a pathogenic mosaic gain of chromosome 12p13, global developmental delay, seizures, and recurrent supraventricular tachycardia presented with fever, vomiting, and respiratory distress. She subsequently developed sudden cardiovascular collapse due to profound

hemolysis, with hemoglobin dropping to 2.7 g/dL. Laboratory findings were consistent with hemolysis, including positive direct antiglobulin test for complement and reticulocytosis [4,6]. *Mycoplasma pneumoniae* infection was confirmed by positive IgM serology [2]. The patient required intensive care support, blood transfusions, corticosteroids, and azithromycin therapy, with eventual clinical and hematological recovery [5,7].

Discussion

Cold agglutinin hemolytic anemia associated with *Mycoplasma pneumoniae* is rare but potentially fatal [7,8]. The pathophysiology involves IgM antibodies directed against red blood cell I antigens, leading to complement activation and intravascular or extravascular hemolysis [6,9]. Severe cases may present with cardiovascular instability, acute renal injury, and multiorgan dysfunction, necessitating intensive care support [5,8]. Management focuses on treatment of the underlying infection, avoidance of cold exposure, transfusion support, and immunomodulatory therapy in selected severe cases [7,10–12].

Conclusion

This case highlights that *Mycoplasma pneumoniae* infection can precipitate severe, rapidly progressive autoimmune hemolytic anemia in medically complex pediatric patients. Early diagnosis and prompt multidisciplinary management are essential to prevent fatal outcomes and ensure recovery [7,8].

Conflict of interest

The authors declare no conflicts of interest.

References

- Bărbulescu D, Jianu M, Stanescu C, Negreanu A. (2017). Severe extrapulmonary complications in *Mycoplasma pneumoniae* infection: a case study. *Romanian Journal of Pediatrics*.;66(1):64–67.
- Sharma D, Kanwar M. (2011). *Mycoplasma pneumoniae*: clinical features and management. *Pulm Med*.;2011:479573.
- Gómez-Barrena G, Martín-López M, Martínez-Montiel O, et al. (2021). Autoimmune hemolytic anemia: case review. *An Pediatr (Engl Ed)*.;95(4):293–299.
- Silva C, Neves G, Silva L, et al. (2019). Clinical-epidemiological profile of pediatric patients with autoimmune hemolytic anemia. *Resid Pediatr*.;9(3):233–238.
- Kumaravel K, Jayakanthan S, Vengatesan A, et al. (2022). Severe hemolytic anemia secondary to *Mycoplasma pneumoniae* infection. *J Med Cases*.;13(3):148–151.
- Ma Y, Zhang Y, Wang Y, et al. (2021). Clinical characteristics of *Mycoplasma pneumoniae* infection in children with cold agglutinin disease. *J Clin Lab Anal*.;35(1):e23632.
- Swiecicki M, Hegerova L, Gertz MA. (2013). Cold agglutinin disease: clinical presentation and management. *Mayo Clin Proc*.;88(1):75–83.
- Dhaliwal G, Cornett R, Tierney LM Jr. (2002). *Mycoplasma pneumoniae* infection and cold agglutinin hemolytic anemia. *West J Med*.;176(3):179–180.
- Hancı M, Güneş A, Altan V, Çolak N. (2017). Severe acrocyanosis precipitated by cold agglutinin secondary to infection. *Turk J Pediatr*.;59(2):209–212.
- Al-Shamri MA. (2005). Hemolytic anemia following *Mycoplasma* infection. *Internet J Hematol*.;4(2). Cleveland Clinic. Cold agglutinin disease: symptoms, diagnosis and treatment. Cleveland Clinic. National Organization for Rare Disorders (NORD). Cold agglutinin disease. NORD.



This work is licensed under Creative Commons Attribution 4.0 License

To Submit Your Article Click Here:

[Submit Manuscript](#)

DOI:10.31579/2690-4861/1001

Ready to submit your research? Choose Auctores and benefit from:

- fast, convenient online submission
- rigorous peer review by experienced research in your field
- rapid publication on acceptance
- authors retain copyrights
- unique DOI for all articles
- immediate, unrestricted online access

At Auctores, research is always in progress.

Learn more <https://auctoresonline.org/journals/international-journal-of-clinical-case-reports-and-reviews>