

History of Resting Tachycardia and Elevated Blood Pressure Episodes in Female Students of Grodno State Medical University

Lelevich A.V *, Cai M., Yankovskaya E.A., Satsuta P.P., Yakimovich E.P., Bon E.I., Malenouskaya M.Y., Misik V.A., Kipen K.M.

Grodno State Medical University.

***Corresponding Author:** Lelevich A.V, Grodno State Medical University.

Received Date: April 06, 2026; **Accepted Date:** April 23, 2026; **Published Date:** May 08, 2026

Citation: Lelevich A.V., Cai M., Yankovskaya E.A., Satsuta P.P., Yakimovich E.P., et al, (2026), History of Resting Tachycardia and Elevated Blood Pressure Episodes in Female Students of Grodno State Medical University, *J, Surgical Case Reports and Images*, 9(3); DOI:10.31579/2690-1897/314

Copyright: © 2026, Lelevich A.V. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract

Arterial hypertension (AH) remains one of the most significant medico-social problems of our time. AH is no longer confined to the elderly population. Modern epidemiological trends indicate a marked 'rejuvenation' of the condition, as recent data point to a growing incidence of high blood pressure among younger individuals, posing a substantial challenge to public health. Studies indicate that a high heart rate (HR) serves not only as a symptom but also as an independent predictor of the development of AH.

Keywords: Arterial hypertension; high heart rate

Introduction

Arterial hypertension (AH) remains one of the most significant medico-social problems of our time. AH is no longer confined to the elderly population. Modern epidemiological trends indicate a marked 'rejuvenation' of the condition, as recent data point to a growing incidence of high blood pressure among younger individuals, posing a substantial challenge to public health. [1]. Studies indicate that a high heart rate (HR) serves not only as a symptom but also as an independent predictor of the development of AH [2]. In individuals with a HR exceeding 80 bpm, the risk of developing sustained hypertension is 2-3 times higher compared to those with normal resting heart rate [3, 4]. This problem is particularly relevant in the student population, where lifestyle-related stress and psycho-emotional strain provoke autonomic imbalance [5].

An increase in HR directly raises cardiac output and shortens the time for diastolic vascular relaxation, creating excessive pressure on the vessel wall [3]. Excess catecholamines simultaneously induce tachycardia and vasoconstriction, increasing total peripheral resistance. A high pulse wave frequency damages the endothelium, reducing nitric oxide synthesis and disrupting vasodilation mechanisms [6]. Sympathetic stimulation activates the renin-angiotensin-aldosterone system, leading to sodium retention, and an increase in circulating blood volume [2].

In the current stratification of cardiovascular risk, special attention is paid to threshold HR values. According to recent clinical guidelines (2024 European Society of Hypertension), a resting HR exceeding 80 bpm is recognized as an independent risk factor for the development of arterial hypertension, as it serves as a clinical marker of sustained sympathetic

overactivity. A HR exceeding 100 bpm is the classic criterion for tachycardia. It has been shown that at such a frequency, the pulse wave becomes "aggressive," causing mechanical trauma to the arterial intima (shear stress), which irreversibly reduces nitric oxide synthesis [6].

Objective

To study the prevalence of episodes of tachycardia at rest and their association with self-reported episodes of elevated/reduced blood pressure Among third-year female students of Grodno State Medical University during the 2025/2026 academic year.

Materials and Methods

The study involved 385 third-year female students at Grodno State Medical University (GRSMU) during the 2025–2026 academic year. Voluntary informed consent was obtained from all participants. The students were anonymously surveyed on the occurrence of tachycardia episodes (exceeding 80 and 100 bpm), as well as self-reported episodes of elevated or reduced blood pressure (response options: "yes," "no," "don't know"). Students who answered "don't know" were excluded from the analysis.

Data were presented as absolute and relative frequencies (%). Comparisons of distributions were performed using Fisher's exact test. The threshold for statistical significance was set at 0.05. Statistical data processing was carried out using StatSoft STATISTICA 10.0 software.

Results and Discussion

It was found that among female students experiencing episodes of resting tachycardia >80 bpm (n=130), episodes of elevated BP were significantly

more frequent compared to those without such episodes (n=180): 31.54% and 17.22%, respectively (p=0.003, Table 1).

History of tachycardia (>80 bpm)	n	No history of elevated BP	History of elevated BP
No	180	82.78% (149)	17.22% (31)
Yes	130	68.46% (89)	31.54% (41)

Table 1: History of tachycardia (>80 bpm) and self-reported episodes of elevated blood pressure in third-year female students at GRSMU, 2025/2026 academic year, % (n)

Among students experiencing episodes of resting tachycardia >100 bpm (n=39), episodes of elevated BP were also more frequent compared to those without such episodes (n=262): 43.59% and 19.47%, respectively (p=0.001, Table 2).

History of tachycardia (>100 bpm)	n	No history of elevated BP	History of elevated BP
No	262	80.53% (211)	19.47% (51)
Yes	39	56.41% (22)	43.59% (17)

Table 2: History of tachycardia (>100 bpm) and self-reported episodes of elevated blood pressure in third-year female students at GRSMU, 2025/2026 academic year, % (n)

When examining the prevalence of reduced BP, it was found that among students with a history of tachycardia >80 bpm (n=138), episodes of hypotension were more frequent compared to those without such episodes (n=181): 65.22% and 43.09%, respectively (p<0.001, Table 3).

History of tachycardia (>80 bpm)	n	No history of hypotension	History of hypotension
No	181	56.91% (103)	43.09% (78)
Yes	138	34.78% (48)	65.22% (90)

Table 3: History of tachycardia (>80 bpm) and self-reported episodes of low blood pressure in third-year female students at GRSMU, 2025/2026 academic year, % (n)

Among students with a history of tachycardia >100 bpm (n=46), episodes of hypotension were also more frequent compared to those without such episodes (n=262): 73.91% and 46.56%, respectively (p<0.001, Table 4).

History of tachycardia (>100 bpm)	n	No history of hypotension	History of hypotension
No	262	53.44% (140)	46.56% (122)
Yes	46	26.09% (12)	73.91% (34)

Table 4: History of tachycardia (>100 bpm) and self-reported episodes of low blood pressure in third-year female students at GRSMU, 2025/2026 academic year, % (n)

The obtained data reflect a significant autonomic imbalance in female students experiencing episodes of resting tachycardia, characterized by increased variability in blood pressure regulation.

Conclusion

The presence of resting tachycardia episodes (>80 and >100 bpm) is associated with increased blood pressure variability, manifested by a significantly higher frequency of both elevated and reduced blood pressure episodes among third-year female students at GrSMU.

References

1. Daniyarova, A. Arterial hypertension in young people: efficacy of sympatholytic therapy / A. Daniyarova, B. Kenenbaeva // Sciences of Europe. – 2022. – Iss. 95. – P. 12–15.
2. 2024 European Society of Hypertension clinical practice guidelines for the management of arterial hypertension / R. Kreutz, M. Brunström, M. Burnier [et al.] // European Journal of Internal Medicine. – 2024. – Vol. 126. – P. 1–15.
3. Rigon, P. A. Activité physique et hypertension: recommandations et pratique clinique [Physical activity and hypertension: guidelines and clinical practice] / P. A. Rigon, G. Wuerzner // Revue Médicale Suisse. – 2025. – Vol. 21, № 930. – P. 1599–1602.
4. Palatini, P. Resting Heart Rate as a Cardiovascular Risk Factor in Hypertensive Patients: An Update / P. Palatini // American Journal of Hypertension. – 2021. – Vol. 34, № 4. – P. 307–317.
5. Stress and Heart Rate Variability during University Final Examination among Lebanese Students / S. Hammoud, R. Karam, R. Mourad [et al.] // Behavioral Sciences. – 2018. – Vol. 9, № 1. – Art. 3.
6. Grassi, G. The Sympathetic Nervous System in Hypertension: Roadmap Update of a Long Journey / G. Grassi // American Journal of Hypertension. – 2021. – Vol. 34, № 12. – P. 1247–1254.



This work is licensed under Creative Commons Attribution 4.0 License

To Submit Your Article Click Here:

Submit Manuscript

DOI:[10.31579/2690-1897/314](https://doi.org/10.31579/2690-1897/314)

Ready to submit your research? Choose Auctores and benefit from:

- fast, convenient online submission
- rigorous peer review by experienced research in your field
- rapid publication on acceptance
- authors retain copyrights
- unique DOI for all articles
- immediate, unrestricted online access

At Auctores, research is always in progress.

Learn more <https://auctoresonline.org/journals/journal-of-surgical-case-reports-and-images>